

**UCC Library and UCC researchers have made this item openly available.
 Please [let us know](#) how this has helped you. Thanks!**

Title	Monocyte mobilisation, microbiota & mental illness
Author(s)	van de Wouw, Marcel; Boehme, Marcus; Dinan, Timothy G.; Cryan, John F.
Publication date	2019-07-19
Original citation	van de Wouw, M., Boehme, M., Dinan, T. G. and Cryan, J. F. (2019) 'Monocyte mobilisation, microbiota & mental illness', Brain, Behavior, and Immunity, 81, pp. 74-91. doi: 10.1016/j.bbi.2019.07.019
Type of publication	Article (peer-reviewed)
Link to publisher's version	http://www.sciencedirect.com/science/article/pii/S0889159119306075 http://dx.doi.org/10.1016/j.bbi.2019.07.019 Access to the full text of the published version may require a subscription.
Rights	© 2019, Elsevier Inc. All rights reserved. This manuscript version is made available under the CC BY-NC-ND 4.0 licence. https://creativecommons.org/licenses/by-nc-nd/4.0/
Embargo information	Access to this article is restricted until 12 months after publication by request of the publisher.
Embargo lift date	2020-07-19
Item downloaded from	http://hdl.handle.net/10468/8570

Downloaded on 2021-11-27T10:55:32Z

Monocyte Mobilisation, Microbiota & Mental Illness

Marcel van de Wouw¹, Marcus Boehme^{1,3}, Timothy G. Dinan^{1,2}, John F. Cryan^{1,3}

¹APC Microbiome Ireland, University College Cork, Cork, Ireland

²Department of Psychiatry and Neurobehavioral Science, University College Cork, Cork, Ireland

³Department of Anatomy and Neuroscience, University College Cork, Cork, Ireland

Corresponding author:

J.F. Cryan: APC Microbiome Ireland; Department of Anatomy and Neuroscience, University College Cork, Cork, Ireland. Email: j.cryan@ucc.ie.

Word count: 6830 excluding reference list, 18014 including reference list.

Keywords: Microbiota; Monocytes; Brain; Behaviour; Immune System; Stress; Ageing; Traumatic Brain Injury; Stroke; Neurodegeneration.

Abstract

The gastrointestinal microbiome has emerged as a key player in regulating brain and behaviour. This has led to the strategy of targeting the gut microbiota to ameliorate disorders of the central nervous system. Understanding the underlying signalling pathways in which the microbiota impacts these disorders is crucial for the development of future therapeutics for improving CNS functionality. One of the major pathways through which the microbiota influences the brain is the immune system, where there is an increasing appreciation for the role of monocyte trafficking in regulating brain homeostasis. In this review, we will shed light on the role of monocyte trafficking as a relay of microbiota signals in conditions where the central nervous system is in disorder, such as stress, peripheral inflammation, ageing, traumatic brain injury, stroke, multiple sclerosis, Alzheimer's disease and Parkinson's disease. We also cover how the gastrointestinal microbiota is implicated in these mental illnesses. In addition, we aim to discuss how the monocyte system can be modulated by the gut microbiota to mitigate disorders of the central nervous system, which will lead to novel microbiota-targeted strategies.

1. Introduction

There is a growing recognition of the role of the gastrointestinal microbiota in regulating brain and behaviour, which has been coined the microbiota-gut-brain axis (Rhee et al. 2009, Collins et al. 2012, Cryan et al. 2012, Cryan et al. 2019). It is crucial to understand the signalling pathways underlying this axis in order to inform future therapeutics in improving CNS functionality in both health and disease. Some pathways have already been widely described, such as host immunity (Rea et al. 2016, Rieder et al. 2017), the vagus nerve (Bonaz et al. 2018, Fulling et al. 2019), metabolite production (Rooks et al. 2016), and enteroendocrine signalling (Plovier et al. 2017, van de Wouw et al. 2017, Lach et al. 2018). In tandem, there is a growing appreciation of the role of monocyte trafficking in regulating brain homeostasis and aiding the brain's response to insult or injury (Ramirez et al. 2017). In this review, we will shed more light on the role of monocyte trafficking as a relay of microbiota signals to the host CNS and its behavioural implications. In addition, we cover how this system can be modified using various microbiota-targeted strategies.

2. Monocytes: from Bone to Brain

Monocytes are mononuclear phagocytes that play crucial roles in tissue homeostasis and immunity, where they are especially important in inflammation and pathogen challenge. Monocytes, in particular those characterised by a high LY6C expression (i.e., LY6C^{hi}), originate in the bone marrow, after which they enter the peripheral circulation. These cells subsequently either differentiate into LY6C^{low} monocytes (Varol et al. 2007, Yona et al. 2013), or differentiate into dendritic cells or macrophages upon entering various tissues (**Figure 1**) (Ginhoux et al. 2014). Notably, specific dendritic cells and most tissue-resident macrophages (e.g., brain microglia) are maintained independently of peripheral monocyte contribution (Ajami et al. 2007, Liu et al. 2009, Varol et al. 2009, Hashimoto et al. 2013, Yona et al. 2013). For an overview of all circulating murine and human subsets, see **Table 1** (for further reading, see (Jakubzick et al. 2017, Guilliams et al. 2018)).




Figure 1. The production and fate of monocytes. Monocytes are produced in the bone marrow through myelopoiesis. LY6C^{high} monocytes subsequently enter the peripheral circulation, after which they either differentiate into LY6C^{low} monocytes, or traffic into other tissues and differentiate in dendritic cells or macrophages. LY6C^{high} monocytes have a high expression of the CCR2 and low expression of the CX3CR1 receptor. LY6C^{low} monocytes have a low expression of the CCR2 and high expression of the CX3CR1 receptor.

Table 1. Monocyte subsets and functions in mice and humans.

Species	Subsets	Surface markers	Chemokine receptors	Function	References
Mouse	LY6C ^{hi}	CD11b ⁺ , LY6C ^{hi}	CCR2 ^{hi} , CX3CR1 ^{Low}	Phagocytosis & Pro-inflammatory	(Palframan et al. 2001, Geissmann et al. 2003, Auffray et al. 2007, Ziegler-Heitbrock et al. 2010)
	LY6C ^{mid}	CD11b ⁺ , LY6C ^{mid}	CCR2 ^{hi} , CX3CR1 ^{Low}	Pro-inflammatory	
	LY6C ^{low}	CD11b ⁺ , LY6C ^{low}	CCR2 ^{low} , CX3CR1 ^{hi}	Patrolling; tissue repair	
Human	Classical	CD14 ⁺⁺ , CD16 ⁻	CCR2 ^{hi} , CX3CR1 ^{Low}	Phagocytosis	(Passlick et al. 1989, Ziegler-Heitbrock et al. 2010, Ziegler-Heitbrock et al. 2013, Patel et al. 2017)
	Intermediate	CD14 ⁺⁺ , CD16 ⁺	CCR2 ^{mid} , CX3CR1 ^{hi} , CCR5 ⁺	Pro-inflammatory	
	Non-classical	CD14 ⁺ , CD16 ⁺⁺	CCR2 ^{low} , CX3CR1 ^{hi} ,	Patrolling	

Infiltration of monocytes into the CNS occurs at a steady-state under non-pathological conditions (Korin et al. 2017, Mrdjen et al. 2018), which has been implicated in brain plasticity (Mohle et al. 2016). Enhanced monocyte trafficking often occurs during various pathological conditions where monocytes can facilitate tissue injury, repair or protection (Ransohoff et al. 2010, Prinz et al. 2011, Ransohoff 2011, Karlmark et al. 2012). Infiltrating monocytes/macrophages exhibit a molecularly and functionally distinct signature compared to the most prominent myeloid cells present in the brain - the microglia (Ginhoux et al. 2010, Hickman et al. 2013, Butovsky et al. 2014, Cronk et al. 2018, Lund et al. 2018). Enhanced cell trafficking can occur through the choroid plexus-cerebrospinal fluid route in many conditions such as stroke (Ge et al. 2017), and traumatic brain injury (TBI) (Szmydynger-Chodobska et al. 2012). In addition, many conditions associated with enhanced cell trafficking into the CNS are often paired with an increased blood-brain barrier permeability, likely allowing monocyte to more easily infiltrate into brain tissue (Takeshita et al. 2012). More recently, it has been shown that myeloid cells can migrate to the brain through vascular channels from the bone marrow in conditions of stroke and aseptic meningitis (Herisson et al. 2018). Under conditions of neuroinflammation, activated microglia release the chemokine C-C motif chemokine ligand 2 (CCL2) and the chemokine (C-X3-C motif) ligand 1 (CX3CL1) (Prinz et al. 2010). The receptors for these ligands, CCR2 and CX3CR1, are expressed on monocytes and subsequently facilitate monocyte trafficking into the brain (Prinz et al. 2010). Whether monocytes play a causal role in the symptomology in specific conditions is often investigated using CCR2-KO or CX3CR1-KO mice, in which monocyte trafficking (into the CNS) is almost ablated (Tsou et al. 2007, Prinz et al. 2010). It is also interesting to note that the CXCR7 receptor has recently been implicated in classical monocyte trafficking into the brain in relation to HIV-associated neurocognitive disorders in humans (Veenstra et al. 2017). Moreover, CCR7 can facilitate TLR4-induced trans-migration of monocytes across the blood-brain barrier *in vitro* (Paradis et al. 2016).

3. Gut microbial shaping of the monocyte system and monocyte function

3.1 The Microbiota in Monocyte Trafficking

Proof-of-principle studies using mice devoid of any microbiota (i.e., germ-free – GF) have revealed that the microbiota is essential for hematopoiesis, as these mice show decreased levels of monocytes in the bone marrow, spleen, and circulation (Balmer et al. 2014, Khosravi et al. 2014). These deficits are ameliorated upon recolonisation of the microbiota or by the administration of microbial-associated molecular patterns (MAMPs) (Khosravi et al. 2014). Similar to GF mice, depletion of the gastrointestinal microbiota by antibiotic administration decreases systemic and brain LY6C^{hi} monocyte levels (Balmer et al. 2014, Khosravi et al. 2014, Möhle et al. 2016), which is restored upon supplementation with the multispecies probiotic VSL#3, which contains eight bacterial strains: *Bifidobacterium breve*, *B. infantis*, *B. longum*, *Lactobacillus acidophilus*, *L. delbrueckii*, *L. paracasei*, *L. plantarum* and *Streptococcus thermophilus* (Möhle et al. 2016). Interestingly, this study also showed that microbiota-regulated monocyte trafficking is essential for healthy adult hippocampal neurogenesis and memory retention (Möhle et al. 2016). Off note, GF mice also show deficits in microglia maturation and function, which is paired with an increase in macrophage colony-stimulating factor 1 receptor expression (CSF1R) on microglia, a key factor in neuron-microglia interactions, indicating a possible role for the gut microbiota in regulating neuron-microglia cross-talk (Erny et al. 2015, Wohleb 2016). Administration of the endotoxin lipopolysaccharide (LPS), a microbiota-derived metabolite, increases monocyte trafficking into the CNS and sickness behaviour (Harden et al. 2015, Rosadini et al. 2017), indicating a role for the microbiota in monocyte-associated neuroinflammation and subsequent behavioural impairment. It is also interesting to note that monocytes can contribute to pathogen dissemination into the brain by carrying microbiota-derived bacterial strains resulting in brain infection (Drevets et al. 2004, Santiago-Tirado et al. 2017).

3.2 Microbial Metabolites and Epigenetic Programming

There is a growing emphasis on the role of microbial metabolites as epigenetic regulators (Stilling et al. 2016, Woo et al. 2017). This is important in regard to monocyte trafficking as prior to their infiltration into tissue and subsequent differentiation into macrophages, monocytes can undergo a trained innate immunity via epigenetic reprogramming, which subsequently determines the function of the cells they differentiate into (Bordon 2014, Guillems et al. 2018). Innate memory training of monocytes has been implicated in the susceptibility to secondary infections and the functional fate of macrophages (Quintin et al. 2012, Bordon 2014, Cheng et al. 2014), and has even been linked to the pathogenesis of neuroimmune psychiatric disorders (Salam et al. 2018). Various microbiota-derived metabolites have already been shown to facilitate trained innate immunity, such as β -glucan, LPS, and butyrate (Saeed et al. 2014, Schulthess et al. 2019). Such data indicates the possibility for the development of microbiota-targeted strategies to reprogram monocytes before their infiltration into the brain to influence the brain and behaviour in healthy and pathogenic conditions (**Figure 2**).




Figure 2: Pathways in which the microbiota can influence monocyte trafficking to the brain and subsequent signalling. A depleted gastrointestinal microbiota results in a decreased myelopoiesis and monocyte levels, resulting in reduced monocyte trafficking to the brain. Alternatively, gut microbiota-derived metabolites, such as lipopolysaccharide (LPS), serotonin (5-HT), SCFAs (short-chain fatty acids), and β-glucans, can influence monocyte properties and the macrophages they subsequently differentiate into.

Perhaps one of the most investigated microbial-derived metabolites is the endotoxin LPS, which is present on the outer membrane of Gram-negative bacteria. LPS activates the toll-like receptor 4 (TLR4) and its co-receptors, CD14 and myeloid differentiation protein 2, resulting in the secretion of various inflammatory cytokines (Rossol et al. 2011, Cavaillon 2018). Incubation of human monocytes with LPS *ex vivo* reprograms monocytes to differentiate into macrophages (i.e., trained immunity) secreting less pro-inflammatory cytokines, such as TNF- α and IL-6 (Saeed et al. 2014).

β -glucans are polysaccharides naturally occurring in cell walls of plants, bacteria, and fungi. The direct effects of β -glucans on immune cells are most likely mediated by cell-surface receptors such as dectin-1 and TLRs, as β -glucans cannot penetrate the cell membrane (Kim et al. 2011). Similar to LPS, β -glucans have also been shown to elicit trained innate immunity (Cheng et al. 2014, Saeed et al. 2014). Specifically, incubation of human monocytes with β -glucan *ex vivo* can reprogram monocytes to differentiate into macrophages secreting more pro-inflammatory mediators, such as TNF- α and IL-6, upon subsequent stimulation with a TLR2 receptor ligand (Cheng et al. 2014, Saeed et al. 2014). It is important to note however, that the immunomodulatory effect of β -glucans is primarily dependent on the source they have been extracted from (for review see: (Sima et al. 2018)).

Another class of gut microbial-derived metabolites known to affect monocyte function are short-chain fatty acids (SCFAs) (for a comprehensive review on SCFAs and host physiology, see: (Koh et al. 2016, Stilling et al. 2016)). The primary source of SCFAs is the microbial fermentation of host-indigestible dietary fibres in the cecum and colon (Macfarlane et al. 2003). The three principle SCFAs are acetate, propionate and butyrate, of which only acetate reaches the peripheral circulation in high concentrations (100–200 μ mol/L, 1- to 15- μ mol/L for propionate and butyrate) (Cummings et al. 1987, Peters et al. 1992, Bloemen et al. 2009). Data indicate an anti-inflammatory effect of SCFA on monocytes, as well as other immune cell populations (Li et al. 2018). All three principle SCFAs can activate the free fatty acid receptor 2 and 3 (FFAR2 and FFAR3 respectively), which are expressed on murine and human monocytes and suppress LPS-induced cytokine secretion (Ang et al. 2016).

Similarly, butyrate inhibits *Staphylococcus aureus*-induced IL-12p40, IL-12p70, and TNF- α secretion, while enhancing IL-10 secretion in human monocytes (Saemann et al. 2000). In addition, human monocytes treated with butyrate have a decreased expression of the innate response receptors CD14 and CD16, which was mediated through the downregulation of the transcription factor PU.1 (Lasitschka et al. 2017), which has been implicated in monocyte/macrophage differentiation (Shen et al. 2016). Indeed, it has recently been shown that human monocytes incubated with butyrate differentiate into macrophages with potent antimicrobial function (Schulthess et al. 2019). In addition, a 4-week oral supplementation of butyrate to individuals with metabolic syndrome influenced subsequent *ex vivo* trained immunity by oxidised low-density lipoprotein, resulting in decreased LPS-induced IL-6 secretion and Pam3CSK4-induced TNF- α secretion, as well as a decreased Pam3CSK4-induced IL-10 production when monocytes were trained with β -glucan (Cleophas et al. 2019).

One of the most studied neurotransmitter systems regulated by the gut microbiota and its metabolites is serotonin (5-HT) (Clarke et al. 2013, Yano et al. 2015). A variety of 5-HT receptors are expressed on monocytes such as the 5-HT_{1A}, 5-HT_{1E}, 5-HT_{2A}, 5-HT_{3A}, 5-HT₄ and 5-HT₇ receptor (Herr et al. 2017). In addition, monocytes express the serotonin transporter and the key enzymes involved in serotonin synthesis (tryptophan hydroxylase - TPH) and for serotonin degradation (monoamine oxidase- MAO) (Herr et al. 2017). Serotonergic signalling in monocytes increases the LPS-induced secretion of the inflammatory cytokines IL-1 β , IL-6, IL-12p40, and IL-8, yet decreases TNF- α (Arzt et al. 1991, Cloez-Tayarani et al. 2003, Durk et al. 2005), highlighting a complex modulatory effect of 5-HT on monocytes. Conversely, 5-HT is also able to influence monocyte to macrophage differentiation, resulting in a decreased LPS-induced inflammatory cytokine release (de las Casas-Engel et al. 2013), potentially showing a role for 5-HT in trained innate immunity.

4. Gut microbiota-targeted therapeutics

There are at many ways by which the gut microbiota can be targeted (long-Smith et al. 2019). For instance, by administrating live bacterial strains that confer health benefits (i.e. probiotics) (Hill et al.

2014, Derrien et al. 2015). Alternatively, administration of host-indigestible dietary fibers, which undergo bacterial fermentation, subsequently stimulates the growth of certain types of bacteria (i.e. prebiotics) (Bindels et al. 2015, Gibson et al. 2017, Stephen et al. 2017). There is also an increased emphasis on synbiotic (i.e. combining both pre- and probiotics) and post-biotics (i.e. microbial-derived metabolites) approaches (Markowiak et al. 2017, Aguilar-Toalá et al. 2018), as well as fermented foods (Kim et al. 2016). If these various microbiota-targeted therapeutics confer mental health benefits, then they are also considered psychobiotics (Dinan et al. 2013, Sarkar et al. 2016). Transplanting the faecal microbiota is also becoming more widely used (Ridaura et al. 2013, Smits et al. 2013, Kelly et al. 2016, Bakker et al. 2017, Kang et al. 2019). Finally, there are also various strategies for depleting specific gut microbes, such as antibiotics (Ianiro et al. 2016, Langdon et al. 2016), and bacteriophages (Dalmasso et al. 2014).

It is also important to note that nutritional interventions can modulate the gut microbiota (Sandhu et al. 2017). For instance, omega-3 fatty acids have been shown to modulate the murine (Pusceddu et al. 2015, Robertson et al. 2017, Provensi et al. 2019), and human gut microbiota (Costantini et al. 2017). In addition, various polyphenols can modulate the composition of the gut microbiota in both rodents (Chen et al. 2016, Sung et al. 2017, Zhou et al. 2018, Campbell et al. 2019), and humans (Most et al. 2017). Similarly, other micronutrients such as vitamin D can modulate the host microbiota (Luthold et al. 2017).

5. Conditions Linked to Increased Monocyte Trafficking

5.1 Acute and Chronic Stress

Stress can be defined as a complex condition wherein the homeostasis of an organism is disturbed (McEwen et al. 2015). This results in an internal response of the organism governed by the hypothalamic-pituitary-adrenal axis (HPA-axis), where the hypothalamus secretes corticotropin-releasing hormone (CRH), stimulating the pituitary gland to secrete adrenocorticotrophic hormone (ACTH), resulting in a secretion of cortisol (corticosterone for rodents) by the adrenal glands (Uchold-

Schmidt et al. 2012, Uschold-Schmidt et al. 2013). Activation of the HPA-axis results in altered levels of glucocorticoids and mineralocorticoids aimed to change the behaviour of the organism. This system becomes hyperactivated upon repeated acute stressors (i.e. chronic stress) resulting in decreased evening corticosterone, a hypersensitive HPA-axis and delayed HPA-axis recovery (Bhatnagar et al. 2006, Finger et al. 2011, Wagner et al. 2011, Uschold-Schmidt et al. 2012, Fuchsl et al. 2013, Iniguez et al. 2014, Langgartner et al. 2016, Peterlik et al. 2017). The crucial role of the HPA-axis within the development of stress-related disorders is highlighted by mouse studies demonstrating an amelioration of chronic stress-induced impairments by the blockade of HPA-axis-related mechanisms, such as corticotropin-releasing factor-2 receptor blockade and adrenalectomy (Cooper et al. 2010, Niraula et al. 2018).

Acute stress and HPA-axis activation are paired with an activation of the immune system as measured by a systemic release of cytokines, and by an increased secretion of cytokines upon immune cell stimulation with LPS and phytohaemagglutinin *ex vivo* (Marsland et al. 2017). However, the elevation of glucocorticoids upon HPA-axis activation is mostly implicated in suppressing the further inflammatory response to an acute stressor (Sapolsky et al. 2000). Glucocorticoids are therefore well-known for their anti-inflammatory effects as determined by their ability to inhibit the expression and action of most cytokines (Brattsand et al. 1996, Coutinho et al. 2011). It is not surprising that they have been the cornerstone for treating inflammatory disease for over 60 years (Hench et al. 1950). However, it is important to note that both acute stress and glucocorticoid administration prior to an LPS challenge potentiate inflammation (Johnson et al. 2002, Frank et al. 2010, Sorrells et al. 2013, Kelly et al. 2018), indicating that glucocorticoids can facilitate both the priming and resolving of inflammation (Sorrells et al. 2010).

Rodent studies exploring how stress influences the immune system have revealed decreased levels of circulating monocyte levels in response to acute stress (Dhabhar et al. 1994), indicating an enhanced monocyte trafficking (Dhabhar et al. 2012). Similarly, cold water-induced stress decreases circulating

monocyte levels in healthy male volunteers (Brazaitis et al. 2014). Stress-induced monocyte trafficking is largely mediated by glucocorticoids, as stress-induced monocyte trafficking is absent in adrenalectomised animals in the active-phase, and is only partially present in the inactive-phase of the day (Dhabhar et al. 1995). In addition, corticosterone administration induces monocyte trafficking in rodents (Dhabhar et al. 2012), while glucocorticoids decrease peripheral monocyte levels in healthy human volunteers (Fauci et al. 1974, Steer et al. 1998, Olnes et al. 2016, Yeager et al. 2016). It is, to our knowledge, still unclear whether monocytes traffic into the brain response to an acute stressor. Even though it is interesting to note that acute stress does induce leukocyte trafficking into the skin (Dhabhar et al. 1996). No differences were found in CNS monocyte levels 14-hours post-stress (Wohleb et al. 2013), even though these acute stress-induced effects might have dissipated at this time point, considering that peripheral monocyte levels are back to baseline 3 hours after an acute stressor (Dhabhar et al. 1995). Notably, there is an increased gene expression of adhesion factors (i.e., ICAM1, VCAM1, E-selectin) and chemokines (i.e., CXCL1 and CXCL2) 14-hours post-stress (Sawicki et al. 2015), which might indicate an enhanced sensitivity for monocytes to traffic and adhere to the CNS in repeated episodes of acute stress.

In line with the fact that acute stress results in activation of the immune system (Marsland et al. 2017), repeated stressors result in sustained low-grade inflammation (Hansel et al. 2010, Rohleder 2014, Langgartner et al. 2018, Niraula et al. 2018). One of the key aspects of this low-grade inflammation is monocytosis as indicated by an increased production of monocytes in the bone marrow (Heidt et al. 2014), and increased levels of circulating, splenic and brain monocytes (Engler et al. 2004, Engler et al. 2005, Engler et al. 2008, Hanke et al. 2012, Wohleb et al. 2014, Sawicki et al. 2015, McKim et al. 2016, Lafuse et al. 2017, Lisboa et al. 2018, McKim et al. 2018, Nie et al. 2018, Gururajan et al. 2019, Yin et al. 2019). Similarly, individuals with a low socioeconomic status, akin to chronic social stress, have increased levels of circulating monocytes (Powell et al. 2013). Monocyte trafficking into the brain is further supported by an increase in blood-brain-barrier permeability in response to chronic stress (Rodriguez-Arias et al. 2015, Lehmann et al. 2018).

The mechanistic role for the monocyte system in the aetiology of chronic stress disorders is indicated by rodent studies utilising knock-out models of CCR2-CCL2 signalling, which is essential for monocyte infiltration into the CNS (Prinz et al. 2010). These studies have revealed that monocyte trafficking mediates chronic stress-induced anxiety-like behaviour (Wohleb et al. 2013). Similarly, chemical dampening of Ly6C^{hi} monocytes in chronically stressed mice ameliorates anhedonia, anxiety-like behaviour and markers of neuroinflammation in the hippocampus, hypothalamus and cortex (Zheng et al. 2016). It is important to note, however, that one study reported the absence of CCR2+ monocyte trafficking into the brain in response to social defeat stress, and that CCR2-KO mice show no alterations in stress-induced microglia activation and depressive-like behaviour (Lehmann et al. 2016).

Glucocorticoids also affect the function of monocytes themselves, which is not surprising considering the expression of glucocorticoid receptors on monocytes (Miller et al. 1998). Similar to the decrease in circulating monocyte levels observed in response to an acute stressor (Dhabhar et al. 2012), murine monocyte treated with glucocorticoids *ex vivo* have an increased migratory and chemotactic activity (Ehrchen et al. 2007). Specifically, glucocorticoid treated human monocytes upregulate the expression of the trafficking receptor CCR2 and have enhanced CCL2-induced chemotaxis (Penton-Rol et al. 1999, Okutsu et al. 2008, Yeager et al. 2016). Functionally, glucocorticoids suppress cytokine secretion upon an immune stimulus, which has been linked to the transcription factors NF- κ B and AP-1 (Coutinho et al. 2011). This is supported by the fact that murine monocytes treated with glucocorticoids have increased phagocytic activity, decreased oxidative burst and apoptosis, and modify T cell responses in an anti-inflammatory manner (Ehrchen et al. 2007, Varga et al. 2014). In fact, glucocorticoids induce the differentiation of monocytes into an activated, anti-inflammatory subset (Varga et al. 2008, Tsianakas et al. 2012). Interestingly, chronic stress is associated with a decreased responsiveness of immune cells to glucocorticoids (Engler et al. 2004, Bailey et al. 2007, Engler et al. 2008, Niraula et al. 2018), which has been coined glucocorticoid resistance (Miller et al. 2002). Importantly, the development of psychosocial stress-induced glucocorticoid resistance in mice is dependent on physical injury (Foertsch et al. 2017).

5.1.1 The Microbiota in Acute and Chronic Stress

Proof-of-principle studies using mice devoid of any microbiota (i.e. germ-free, GF) have demonstrated a role for the microbiota in stress responsiveness (Sudo et al. 2004, Vodicka et al. 2018), and stress-related receptor pathway genes in the hippocampus (Luo et al. 2018). This postulates the rationale for modulating acute stress-responses using microbiota-targeted strategies. Indeed, bacterial strains such as *L. rhamnosus* and *B. bifidum* reduce HPA-axis reactivity in rodents (Bravo et al. 2011, Fukui et al. 2018), and a *B. longum* strain in healthy volunteers (Allen et al. 2016). Similarly, the prebiotic sugars fructooligosaccharides (FOS) and galactooligosaccharides (GOS) decreases stress-responsiveness in rodents (Burokas et al. 2017), and a supplement containing FOS and B-GOS decreases cortisol levels in humans (Schmidt et al. 2015). The combination of *L. casei* and inulin (i.e. synbiotics), decreases stress-responsiveness in rats (Barrera-Bugueno et al. 2017). And not surprisingly, administration of the microbial-derived metabolites SCFAs, which are mostly derived from the gut microbial fermentation of dietary fibres, ameliorates stress-responsiveness in mice that had previously undergone psychosocial stress (van de Wouw et al. 2018).

In conditions of chronic stress, alterations in the composition and functional capacity of the gut microbiota have been detected in rodent models (Bailey et al. 2011, Galley et al. 2014, Bharwani et al. 2017, Burokas et al. 2017, Galley et al. 2017, Marin et al. 2017, Szyszkowicz et al. 2017, Dunphy-Doherty et al. 2018). Considering that microbiota-targeted strategies can reduce the response to an acute stressor, it is not surprising that these strategies are also able to reduce chronic stress-induced deficits in behaviour. For instance, supplementation of various bacterial strains from the class *Lactobacillus* results in beneficial effects on chronic stress-induced cognitive impairments, anxiety- and depressive-like behaviour in rodents (Liang et al. 2015, Bharwani et al. 2017, Marin et al. 2017, Dhaliwal et al. 2018, Li et al. 2018). Similarly, the bacterial strains *L. casei* Shirota YIT 9029 and *L. plantarum* 299v reduce cortisol levels in students undergoing exam stress (Andersson et al. 2016, Takada et al. 2016). Prebiotic supplementation in mice undergoing psychosocial stress prevents deficits in stress-responsiveness, anxiety- and depressive-like behaviour in mice (Tarr et al. 2015,

Burokas et al. 2017). Moreover, administration of the microbial-derived metabolites SCFAs ameliorates stress-induced deficits in reward-seeking behaviour and stress-responsiveness (van de Wouw et al. 2018).

Overall, these data show great promise for gut microbiota-targeted nutritional interventions for alleviating stress-related conditions. Overall, the fact that acute stress and glucocorticoids can influence monocyte functionality makes monocytes an interesting target in alleviating stress-related disorders. Nonetheless, more research is warranted on monocyte receptors and (associated) signalling pathways to develop further targeted nutritional interventions for stress-related disorders.

5.2 Peripheral Inflammation, Encephalitis and Sickness Behaviour

Sickness behaviour is a crucial part of the body's mechanisms in the battling of, and recovering from infectious disease (Harden et al. 2015). The core symptoms associated with sickness behaviour are fatigue, social withdrawal, malaise, hyperalgesia, disturbed sleep and cognitive dysfunction, and a loss of appetite and interest in usual daily activities (Harden et al. 2015). These symptoms are often the result of increased pro-inflammatory cytokine signalling in the CNS originating from the periphery (Dantzer et al. 2008). Even though sickness behaviour during infectious disease expedites recovery, enduring immune activation in the brain can exacerbate sickness behaviour and induce the development of depression symptomatology in vulnerable individuals (Dantzer et al. 2008). One aspect of sickness behaviour-associated inflammatory signalling is monocytosis and increased CCR2+ monocyte trafficking into the CNS (Kerfoot et al. 2006, Getts et al. 2008, D'Mello et al. 2009, D'Mello et al. 2013, D'Mello et al. 2015, White et al. 2016, D'Mello et al. 2017, Andonegui et al. 2018, Canivet et al. 2019). The role of monocyte trafficking in sickness behaviour is demonstrated by the fact that CCR2-KO mice do not show any cognitive impairments induced by *Streptococcus pneumoniae* infection (Andonegui et al. 2018). Similarly, blockade of hepatic inflammation-induced monocyte trafficking into the brain using anti-P selectin and anti- $\alpha 4$ integrin attenuates sickness behaviour (D'Mello et al. 2009). This study also showed the necessity of TNF- α signalling in the recruitment of monocytes into the CNS

in mice with hepatic inflammation (D'Mello et al. 2009). It is therefore also interesting to note that peripheral and central TNF- α signalling induces sickness behaviour (Palin et al. 2007, Klaus et al. 2016). Moreover, understanding how monocyte trafficking contributes to such pathways may also have implications for understanding and managing aspects of human fatigue (Dantzer et al. 2014).

It is also interesting to note that, similar to peripheral inflammation, virus infection-induced encephalitis promotes CX3CR1⁺ monocytes trafficking into the brain and TNF- α signalling, which has been implicated in synaptic and learning deficits (Garre et al. 2017). A causal role of monocyte trafficking in the development of neuroinflammation is revealed by the use of CCR2-KO and CX3CR1-KO mice, which show a prolonged survival and prevention of neuroinflammation and neuronal death in response to viral encephalitis, even though seizure development remains unaffected (Kaufer et al. 2018).

5.2.1 The Microbiota in Peripheral Inflammation and Sickness Behaviour

Inflammation-induced anxiety-like behaviour caused by the administration of LPS administration is ameliorated by supplementation of the prebiotic GOS mixed with lactose, glucose and galactose in mice (Savignac et al. 2016). Similarly, bile duct ligation-induced sickness behaviour is ameliorated by administration of the probiotic VSL#3, and reduces monocyte infiltration into the brain and microglia activation (D'Mello et al. 2015). These data indicate a promising role for microbiota-targeted strategies in ameliorating sickness behaviour, potentially through reducing monocyte trafficking.

5.3 Ageing

The immune system undergoes a dynamic change during the process of ageing, which is often characterised by a chronic low-grade inflammation, also termed inflammageing (Franceschi et al. 2000, Oishi et al. 2016). This is associated with a decreased response to various immune stimuli, defined as immunosenescence (van Duin et al. 2007, Shaw et al. 2013, Frasca et al. 2016). In addition, microbial metabolites such as LPS are systemically elevated, potentially due to an increase in intestinal permeability, which may contribute to inflammageing (Kim et al. 2016, Scott et al. 2017). One key

aspect of ageing-associated inflammation is monocytosis and increased myeloid cell levels in the brain (Martinez et al. 2015, Puchta et al. 2016, Martin et al. 2017, Boehme et al. 2019). Furthermore, these monocytes are primed to produce more cytokines in response to LPS (Hearps et al. 2012, Puchta et al. 2016). One critical factor driving age-related monocytosis and dysfunction is the inflammatory cytokine TNF- α (Puchta et al. 2016), which is also able to polarise macrophages into a proinflammatory state (Oishi et al. 2016). Increased circulating and intracellular levels of TNF- α have been reported in monocytes of ageing individuals (Bruunsgaard et al. 1999, Hearps et al. 2012). It is also interesting to note that age-related memory decline is correlated with longitudinal increases in CCL2 levels (Bettcher et al. 2019), further indicating that the monocyte system adapts a more pro-inflammatory phenotype in response to ageing. It is also important to note that ageing is associated with a deterioration of other systems associated with monocyte trafficking, such as the HPA-axis (Vitlic et al. 2014), as well as an increased vulnerability to conditions associated with monocyte trafficking, such as TBI and stroke (Crapser et al. 2016, Sun et al. 2019).

5.3.1 The Microbiota in Ageing

Shifts in the composition of the gut microbiota with age have been detected in rodent models (Fransen et al. 2017, Scott et al. 2017, Thevaranjan et al. 2017, Spychala et al. 2018, van der Lugt et al. 2018, Boehme et al. 2019), and humans (Biagi et al. 2010, Claesson et al. 2012). Targeting the microbiota using a prebiotic-enriched diet ameliorates both monocyte influx and subsequent microglia activation in middle-aged mice (Boehme et al. 2019). This is in line with another study demonstrating that a high fibre diet containing inulin reduces the expression of inflammatory markers in microglia of aged mice (Matt et al. 2018). Similarly, supplementation of the probiotic *L. plantarum* KY1032 and *L. curvatus* HY7601 ameliorates memory impairments in aged rats (Jeong et al. 2015), while *L. helveticus* IDCC3801 improves cognition in aged adults (Chung et al. 2014). Notably, the bacterial strain *L. reuteri* DSM 17938 does not affect measures of depression, anxiety and perceived stress in aged adults (Östlund-Lagerström et al. 2016), potentially indicating that the positive effects of probiotics on age-associated deficits might be bacterial strain dependent.

5.4 Traumatic Brain Injury

The attraction of Ly6C^{hi}-monocytes to the site of the lesion is an early response phenomenon in TBI, resulting in neuroinflammation (Hsieh et al. 2013, Loane et al. 2016, Makinde et al. 2017). The importance of monocyte trafficking within TBI symptomatology is indicated by studies using CCR2-KO mice, as these mice show a decreased lesion volume, decreased pro-inflammatory response and improved cognition following TBI (Hsieh et al. 2014, Israelsson et al. 2014, Gyoneva et al. 2015, Chou et al. 2018). Similarly, CCR2 antagonism results in the downregulation of the pro-inflammatory response and long-term cognitive dysfunction in response to TBI (Morganti et al. 2015). Whereas the cognate ligand of the CCR2 receptor, the chemokine CCL2, is upregulated in the first hours following TBI in both humans (Semple et al. 2010) and rodents (Israelsson et al. 2008, Rhodes et al. 2009). The importance of monocyte trafficking is further indicated by the fact that monocyte depletion attenuates TBI-induced ventricle enlargement and improves neural connectivity, even though lesion size remains unaffected (Makinde et al. 2018). It is interesting to note, however, that CX3CR1-KO mice show reduced sensorimotor deficits and cellular damage four days post-TBI, which was subsequently worsened five weeks post-TBI (Zanier et al. 2016). Such data might indicate that CX3CR1⁺ monocyte trafficking is involved in long-term tissue repair and the resolving of inflammation in TBI. Conversely, CCR2-KO mice show decreased TBI-induced long-term cognitive impairments (Hsieh et al. 2014, Morganti et al. 2015), indicating that long-term impairments might be modulated by monocytes in a subset-specific manner.

5.4.1 The Microbiota in Traumatic Brain Injury

The composition of the gastrointestinal microbiota changes in response to TBI in mice (Treangen et al. 2018). Targeting the microbiota by supplementing *Clostridium butyricum* to mice that underwent TBI results in an amelioration in neurological severity score, brain oedema and neurodegeneration (Li et al. 2018). Similar findings have been reported for *L. acidophilus* showing an ameliorated neurological severity score, neurodegeneration and lesion volume (Ma et al. 2019). Furthermore, *B. longum*, *L. bulgaricus*, and *Streptococcus thermophilus* supplementation shortens the stay in the intensive care

unit of TBI patients (Tan et al. 2011). Which has also been reported for TBI patients receiving a formulation containing (glutamine and) fermented milk with *L. johnsonii* (Falcão de Arruda et al. 2004). Overall, findings suggest some promise in reducing TBI-symptomatology using microbiota-targeted strategies, but more evidence is warranted (Brenner et al. 2017).

5.5 Stroke

Stroke is characterized by a disruption of the blood supply to the CNS leading to focal ischemia, apoptosis, neuroinflammation, and functional decline, and is also associated with increased monocyte trafficking into the CNS (Wattananit et al. 2016, Ge et al. 2017, Weise et al. 2017, Garcia-Bonilla et al. 2018). Trafficking of monocytes into the CNS following a stroke can occur through the choroid plexus-cerebrospinal fluid route (Ge et al. 2017), as well as through the blood-brain barrier, which is more permeable in response to stroke (Moskowitz et al. 2010). These infiltrated monocytes subsequent contribute to the recovery from stroke (Wattananit et al. 2016). In addition, LPS-induced immune system priming enhances monocyte trafficking into the brain and suppresses post-ischemic inflammation following a stroke, which was absent in CCR2-KO mice (Garcia-Bonilla et al. 2018). Similarly, administration of *in vitro* IL-4- and IL-13-primed monocyte before the induction of stroke, ameliorates stroke-induced motor and cognitive dysfunction, which was independent of the infarct size (Ge et al. 2017).

5.5.1 The Microbiota in Stroke

The composition of the gut microbiota changes in response to severe stroke (Singh et al. 2016, Stanley et al. 2016, Stanley et al. 2018, Xia et al. 2019). The role of the microbiota in stroke outcome is highlighted by the findings that GF mice, and mice that were administered additional antibiotics, show a reduced infarct volume in response to stroke (Benakis et al. 2016, Singh et al. 2018). Even though antibiotic treatment has also been reported to worsen stroke outcome in mice (Winek et al. 2016). Not surprisingly, transplantation of the microbiota from mice previously undergone stroke to GF mice results in larger infarct volumes in response to a subsequent stroke (Singh et al. 2016). Whereas the

transfer of a healthy mouse microbiota to mice one-day post-stroke reduces infarct volumes (Singh et al. 2016). Similarly, transplantation of the gut microbiota from acute ischemic stroke patients with a relatively more altered microbiota into mice results in a worse stroke outcome, compared to mice receiving the microbiota from stroke patients with relatively less stroke-induced alterations in gut microbiota composition (Xia et al. 2019). These data indicate a causal role for the microbiota in stroke responsiveness.

5.6 Multiple Sclerosis

Enhanced monocyte trafficking into the CNS has been observed in various conditions of neurodegeneration (Baufeld et al. 2018), such as human multiple sclerosis (MS) (Fischer et al. 2019), and its associated animal model experimental autoimmune encephalomyelitis (EAE) (Fife et al. 2000, Izikson et al. 2000, Mildner et al. 2009, Ajami et al. 2011, Jordao et al. 2019). MS is an autoimmune disease where the immune system attacks nerve cells leading to demyelination and disruption of nerve signalling. In EAE, CCR2⁺ monocytes migrate into the brain as early effector cells and facilitate the development of EAE (Mildner et al. 2009, Ajami et al. 2011), suggesting that monocytes may play a pivotal role during the early phases of MS-associated neuroinflammation and autoimmunity. Indeed, CCR2-KO mice show reduced inflammation and delayed disease onset (Fife et al. 2000, Izikson et al. 2000, Columba-Cabezas et al. 2002, Ajami et al. 2011). It is also interesting to note that monocytes from MS patients treated with type-I interferon, a common treatment for MS, secrete less IL-1 β , a pro-inflammatory cytokine (Guarda et al. 2011).

5.6.1 The Microbiota in Multiple Sclerosis

The composition of the gut microbiota is altered in patients with MS (Chen et al. 2016, Berer et al. 2017, Cekanaviciute et al. 2017, Zeng et al. 2019) and mice with EAE (He et al. 2019). A causal role for the microbiota in MS pathophysiology is illustrated by transferring the microbiota from MS patients to mice, resulting in a fastened EAE progression (Berer et al. 2017, Cekanaviciute et al. 2017). It is also interesting to note that intermittent fasting confers protection to EAE through the gut microbiota

(Cignarella et al. 2018). Similarly, depletion of the microbiota using antibiotics in a mouse model of MS prevents motor dysfunction, limits axon damage and delays the onset of clinical score, while decreasing spatial memory (Mestre et al. 2019, Zeraati et al. 2019). These results indicate a promising prospect of alleviating MS symptomatology using microbiota-targeted strategies. Interestingly, MS patients supplemented with a probiotic containing *Lactobacillus*, *Bifidobacterium* and *Streptococcus* reduced the frequency of classical monocytes (Tankou et al. 2018). Similarly, supplementation with *L. acidophilus*, *L. casei*, *B. bifidum*, and *L. fermentum* improves expanded disability status scale scoring, and depression and anxiety measures (Kouchaki et al. 2017). In addition, *L. reuteri* administration reduces the development of EAE in mice (He et al. 2019).

5.7 Alzheimer's Disease

Alzheimer's disease (AD) is associated with an enhanced myeloid cell trafficking into the brain, compared to normal ageing (Martin et al. 2017). AD is a neurodegenerative disease associated with neuroinflammation, neuronal cell death, an accumulation of amyloid- β in the brain and progressive dementia. Monocytes derived from AD mouse models and late-stage patients with AD show an inflammatory phenotype (Saresella et al. 2014, Martin et al. 2017, Thome et al. 2018). These monocytes seem to be involved in the clearance of amyloid- β (Michaud et al. 2013, Martin et al. 2017, Guo et al. 2019), indicating that these cells can reduce AD pathophysiology. In fact, bone marrow-derived macrophages are more efficacious in amyloid- β phagocytosis and clearance than brain microglia (Simard et al. 2006). The protective role of monocytes in AD is further illustrated by the fact that CCR2-KO accelerates amyloid- β accumulation in the hippocampus and prefrontal cortex and expedites cognitive decline in mouse models of AD (El Khoury et al. 2007, Mildner et al. 2011, Naert et al. 2011, Naert et al. 2012). It is also interesting to note that elevated plasma CCL2 levels have been reported in individuals with early-stage AD, even though CCR2 receptor levels on monocytes were decreased (Zhang et al. 2013).

5.7.1 The Microbiota in Alzheimer's Disease

The composition of the gut microbiota is altered in patients with AD (Unger et al. 2016, Vogt et al. 2017, Zhuang et al. 2018), as well as in rodent models (Harach et al. 2017, Sun et al. 2019). Interestingly, a GF mouse model of AD shows a reduced amyloid- β accumulation (Harach et al. 2017). And similarly, depletion of the gut microbiota using an antibiotic cocktail reduces amyloid- β deposition yet increases soluble A β levels (Minter et al. 2016). Targeting the gut microbiota in a mouse model of AD using a probiotic formulation containing *S. thermophiles*, *B. longum*, *B. breve*, *B. infantis*, *L. acidophilus*, *L. plantarum*, *L. paracasei*, *L. delbrueckii subsp. bulgaricus*, *L. brevis* reduces amyloid- β deposition, brain damage and cognitive decline (Bonfili et al. 2017, Bonfili et al. 2018). Similarly, supplementing a probiotic containing *L. acidophilus*, *L. casei*, *B. bifidum*, and *L. fermentum* improves cognition to AD patients improves cognition (Akbari et al. 2016).

5.8 Parkinson's Disease

Parkinson's disease (PD) is the most common neurodegenerative movement disorder and is characterised by a progressive loss of dopaminergic neurons in the substantia nigra pars compacta along with the presence of Lewy bodies and Lewy neurites composed of α -synuclein throughout the CNS (Harms et al. 2018). PD is also associated with increased infiltration of monocytes into the brain (Harms et al. 2018). Interestingly, CCR2-KO mice show attenuated α -synuclein-induced neuroinflammation and degeneration of dopaminergic neurons (Harms et al. 2018). Functionally, monocytes from patients with PD show enhanced phagocytoses and reduced glucocerebrosidase activity, of which the latter is often associated with PD symptomatology (Atashrazm et al. 2018, Wijeyekoon et al. 2018).

5.8.1 The Microbiota in Parkinson's Disease

Alterations in the composition of the gastrointestinal microbiota have been observed in both PD patients (Hasegawa et al. 2015, Keshavarzian et al. 2015, Scheperjans et al. 2015, Aho et al. 2019, Barichella et al. 2019), and animal models (Sampson et al. 2016). Transferring the microbiota from PD

patients to a GF mouse model of PD results in α -synuclein-mediated motor deficits (Sampson et al. 2016). Furthermore, the GF mouse model of PD shows ameliorated motor and gastrointestinal dysfunction, microglia inflammation and α -synuclein pathology, compared to their conventional PD counterparts (Sampson et al. 2016). Subsequent SCFA supplementation to these GF mice promotes α -synuclein-mediated neuroinflammation (Sampson et al. 2016). Whereas individuals who have undergone a vagotomy, disrupting one of the major pathways in microbiota-gut-brain communication, have reduced risk of developing PD (Svensson et al. 2015, Liu et al. 2017). Importantly, it was recently even shown that propagation of α -synuclein from the gut to the brain via the vagus nerve causes PD in mice (Kim et al. 2019). Overall, these results show a causal role for the gut microbiota in PD pathology. Finally, it is interesting to note that gut microbial tyrosine decarboxylases decrease levels of levodopa in the treatment of PD (van Kessel et al. 2019).

6. Conclusion

It is crucial that we appreciate the complex role of monocyte trafficking in regulating brain function and its behavioural consequences in order to understand their role in brain health and disease. Deficiencies in monocyte trafficking in absence of sickness or injury results in impaired adult hippocampal neurogenesis and memory retention (Mohle et al. 2016), whereas excessive monocyte trafficking is associated with negative consequences in conditions such as viral encephalitis (Getts et al. 2008, Kaufer et al. 2018), TBI (Hsieh et al. 2014, Israelsson et al. 2014), chronic stress (Wohleb et al. 2013, Mackos et al. 2016), EAE (Fife et al. 2000, Izikson et al. 2000, Columba-Cabezas et al. 2002, Ajami et al. 2011) and PD (Harms et al. 2018). Conversely, these infiltrating monocytes also contribute to the recovery from stroke (Wattananit et al. 2016) and mitigate the development of AD (El Khoury et al. 2007, Mildner et al. 2011, Naert et al. 2011, Naert et al. 2012). Thus, whether monocyte infiltration into the brain is beneficial or harmful is condition-dependent, and modulating monocyte trafficking and their properties should therefore be tailored to each disease condition. The gut microbiota could play a pivotal role here (**Figure 3**), as many versatile and distinct microbiota-targeted

strategies have already been developed (Table 2). Even though many of these microbiota-targeted strategies have not been investigated in light of monocyte trafficking, some research already shows possible implications.




Figure 3: The therapeutic potential of microbiota-targeted strategies in conditions associated with increased monocyte trafficking into the brain. The microbiota can (epigenetically) program circulating monocytes, prior to their trafficking into the brain and subsequent differentiation into macrophages. These monocytes and monocyte-derived macrophages play a causal role in specific aspects of pathophysiology of various disease conditions, such as traumatic brain injury (TBI), experimental autoimmune encephalomyelitis (EAE) and human multiple sclerosis (MS), Parkinson's disease (PD), Alzheimer's disease (AD) and stroke, as well as conditions associated with chronic stress and sickness. Monocytes play a negative role (red boxes) or positive role (green boxes) depending on the disease condition.

We have shown that a prebiotic-enriched diet ameliorated both monocyte trafficking into the brain and microglia activation in middle-aged mice (Boehme et al. 2019), which is similar to another study demonstrating that a high fibre diet containing inulin reduces the gene expression of inflammatory markers in the microglia of aged mice (Matt et al. 2018). Conversely, antibiotic administration decreases brain LY6C^{hi} monocyte levels in healthy mice, resulting in deficits in neurogenesis and memory retention, which was subsequently restored upon supplementation with the probiotic VSL#3 (Möhle et al. 2016). And similarly, VSL#3 has also been shown to attenuate the development of sickness behaviour, reduce microglia activation and decrease monocyte infiltration into the brain following bile duct ligation in mice (D'Mello et al. 2015). However, more research is needed to further illuminate microbiome-host interactions in the context of monocytes trafficking to the CNS and their implications in brain health and function, even though initial data shows great promise.

However, there are still many gaps in our understanding of how we can target monocyte trafficking to ameliorate disease symptomatology. Even though we know whether monocytes play a positive or negative role in many of these disease conditions, more information is needed as to how monocytes locally exert their effects in the brain. Novel techniques such as metabolomics and single-cell sequencing will likely play a pivotal role in the dissection of the role of monocytes in impacting the brain microenvironment. Recent studies suggest a therapeutic role for monocytes in AD, as they seem to be involved in the clearance of amyloid- β (Michaud et al. 2013, Martin et al. 2017, Guo et al. 2019). Knowing this allows us to screen for microbiota-targeted therapeutics, such as bacterial strains and metabolites, that enhance the ability of monocytes to clear amyloid- β . Particular in light of the ageing society, microbiota-based strategies that modulate monocyte behaviour could have a tremendous impact on novel therapeutics and the healthcare industry. However, we also need a better understanding of how and whether specific microbiota-derived metabolites can impact monocytes in the circulation prior to attachment to the endothelial vasculature and diapedesis and how this affects the macrophages they differentiate into. This would provide us with a better understanding of how we can modulate monocyte trafficking using microbiota-targeted strategies. Nonetheless, many

advances have been made over the last few years, which provides a promising prospect for microbiota-targeted strategies in altering monocytes and their functionality. Overall, this review provides an exciting starting point for the development of such therapeutics in improving brain physiology and behaviour in both health and disease.

Table 2. A summary of microbiota-targeted strategies studies for placebo-controlled human studies and animal studies.

Species/Model and Sex	Gut Microbiota-Targeted Strategy	Treatment Dosage	Time and	Outcome	References
Multiple sclerosis (M/F)	<i>Lactobacillus acidophilus</i> , <i>L. casei</i> , <i>Bifidobacterium bifidum</i> , and <i>L. fermentum</i>	2×10^9 CFU daily for 12 weeks		Decreased depression and anxiety measures	(Kouchaki et al. 2017)
Individuals (M/F) affected by daily stress during the last month	Probio-Stick: <i>L. acidophilus</i> Rosell-52 and <i>B. longum</i> Rosell-175	3×10^9 CFU per day for 3 weeks		Decrease in specific stress-induced gastrointestinal symptoms	(Diop et al. 2008)
Student (M/F) exam stress	Milk fermented with <i>L. casei</i> Shirota YIT 9029	1×10^9 CFU per day for 8 weeks		Ameliorated cortisol 1 day prior to the exam.	(Takada et al. 2016)
Student (M/F) exam stress	<i>L. plantarum</i> 299v	1×10^{10} CFU per day for 2 weeks		Decreased cortisol	(Andersson et al. 2016)
Brain injury (M/F)	30 g glutamine and 240 ml fermented milk with <i>L. johnsonii</i>	Enteral administration for 6-14 days		Shortened stay in the intensive care unit	(Falcão de Arruda et al. 2004)
Traumatic brain injury (M/F)	<i>B. longum</i> , <i>L. bulgaricus</i> , and <i>Streptococcus thermophilus</i> .	Enteral administration 1×10^9 CFU split over 7 dosages daily for 21 days		Shortened stay in the intensive care unit	(Tan et al. 2011)
Aged adults (M/F) (50-70 years)	Lactononadecapeptide with <i>L. helveticus</i> CM4	2.4 g Lactononadecapeptide with <i>L. helveticus</i> fermented milk per day for 8 weeks		Improved cognition	(Ohsawa et al. 2018)
Aged adults (M/F) (>65 years)	<i>L. reuteri</i> DSM 17938	1×10^8 CFU twice daily for 12 weeks		No significant effect on depression, anxiety or perceived stress	(Östlund-Lagerström et al. 2016)
Aged adults (M/F) (60-75 years)	<i>L. helveticus</i> IDCC3801	Four tablets daily for 12 weeks		Improved cognition	(Chung et al. 2014)
Alzheimer's disease (M/F)	<i>L. acidophilus</i> , <i>L. casei</i> , <i>B. bifidum</i> , and <i>L. fermentum</i>	2×10^9 CFU daily for 12 weeks		Improved cognition	(Akbari et al. 2016)
C57Bl/6 mice (M), LPS-induced sickness-behaviour	B-GOS®: 48% w/w GOS, 26% lactose, 14% glucose and 12% galactose	For 3 weeks in drinking water		Ameliorated anxiety-like behaviour	(Savignac et al. 2016)

C57Bl/6 mice (M), Traumatic brain injury	<i>L. acidophilus</i>	1 × 10 ¹⁰ CFU daily by oral gavage for 1 week starting 4 hours post operation	Decreased neurological severity score, neurodegeneration and lesion volume	(Ma et al. 2019)
C57Bl/6 mice (M), Traumatic brain injury	<i>Clostridium butyricum</i>	1 × 10 ⁹ CFU daily by oral gavage for 2 weeks before traumatic brain injury and for 2 weeks afterwards	Ameliorated neurological severity score, brain oedema and neurodegeneration	(Li et al. 2018)
F344 rats (M), Stress	<i>L. casei</i> Shirota YIT 9029	In food for 2 weeks	Ameliorated acute stress-induced cortisol	(Takada et al. 2016)
F344 rats (M), Stress	GOS, or polydextrose alone, lactoferrin alone, or GOS, polydextrose and lactoferrin	GOS, 7 g/kg; polydextrose, 7 g/kg; lactoferrin, 2.6 g/kg in food for 4 weeks	Ameliorated stress-induced learned helplessness	(Mika et al. 2017)
C57Bl/6 mice (M), Chronic stress	FOS and GOS	0.3-0.4 g/mouse per day in drinking water for 6 weeks	Ameliorated cognitive dysfunction, anhedonia, anxiety-like and depressive-like behaviour	(Burokas et al. 2017)
C57Bl/6 mice (M), Chronic stress	3'Sialyllactose or 6'Sialyllactose	In food (5%) for 2 weeks and during stress (1 week)	Ameliorated anxiety-like behaviour	(Tarr et al. 2015)
C57Bl/6 mice (M), Chronic stress	<i>L. reuteri</i>	2 × 10 ⁹ CFU in food for 4 weeks	Ameliorated despair behaviour	(Marin et al. 2017)
C57Bl/6 mice (M), Chronic stress	<i>L. rhamnosus</i> JB-1	1.67 × 10 ⁹ CFU per dose by oral gavage. 20 dosages over 28 days	Ameliorated social and anxiety-like behaviour	(Bharwani et al. 2017)
C57Bl/6 mice (M), Chronic stress	<i>L. helveticus</i> MCC1848	±1 × 10 ⁹ CFU in food for 2 weeks	Ameliorated anhedonia	(Maehata et al. 2019)
Sprague–Dawley rats (M), Chronic stress	<i>L. helveticus</i> NS8	1 × 10 ⁹ CFU/mL in drinking water for 3 weeks	Ameliorated corticosterone levels, anhedonia and anxiety-like behaviour	(Liang et al. 2015)
Swiss albino LACA mice (M), Chronic stress	<i>L. plantarum</i> MTCC 9510	2 × 10 ¹⁰ CFU daily by oral gavage for 4 weeks	Ameliorated cognitive impairments, anxiety- and depressive-like behaviour	(Dhaliwal et al. 2018)

Kunming mice (M), Chronic stress	<i>L. kefiranofaciens</i> ZW3	1 × 10 ⁸ CFU daily by oral gavage for 6 weeks	Ameliorated depressive-like behaviour	(Sun et al. 2019)
C57Bl/6 mice (M), Chronic stress	Short-chain fatty acids	Sodium acetate (67.5mM), sodium propionate (25mM), and sodium butyrate (40mM) in drinking water for 11 weeks	Ameliorated anhedonia and stress-responsiveness	(van de Wouw et al. 2018)
Sprague–Dawley rats (M), Chronic stress	Short-chain fatty acids	Sodium propionate (200 mmol/L) 1 mL intrarectally daily for 5 weeks.	Ameliorated depressive-like behaviour	(Li et al. 2018)
Fischer 344 Rats (M), aged (18 months)	<i>L. plantarum</i> KY1032 and <i>L. curvatus</i> HY7601	1 × 10 ¹⁰ CFU by oral gavage for 8 weeks	Ameliorated memory impairment	(Jeong et al. 2015)
C57Bl/6 mice (M), middle-aged (10 months)	FOS-inulin	In food (9.2% inulin, 0.8% FOS) for 3.5 or 14 weeks	Ameliorated infiltration of LY6C ^{hi} monocytes into the brain and reduced microglia activation	(Boehme et al. 2019)
Balb/c mice (M), aged (22-25 months)	Inulin	In food (5%) for 4 weeks	Ameliorated expression of genes associated with inflammation in microglia	(Matt et al. 2018).
C57Bl/6 mice injected with myelin oligodendrocyte glycoprotein peptide and pertussis toxin (F), Experimental autoimmune encephalomyelitis	<i>L. reuteri</i>	1 × 10 ¹⁰ CFU by oral gavage for 20 days	Reduced clinical score	(He et al. 2019)
C57Bl/6 mice injected with myelin oligodendrocyte glycoprotein peptide (F), Experimental autoimmune encephalomyelitis	Antibiotic cocktail	neomycin (10 mg/ml), ampicillin (1 mg/ml), metronidazole (10 mg/ml), vancomycin (5 mg/ml) and Amphotericin-B (0.1 mg/ml) via drinking water for 12 weeks	Reduced onset of clinical score and decreased spatial memory	(Zeraati et al. 2019)

TMEV-IDD SJL/J mice (F), Experimental autoimmune encephalomyelitis	Antibiotic cocktail	ampicillin (1 mg/mL), metronidazole (1 mg/mL), neomycin sulfate (1 mg/mL) and vancomycin (0.5 mg/mL) in drinking water for 2 weeks	Reduced motor dysfunction and axon damage	(Mestre et al. 2019)
APP^{SW}E/PS1^{ΔE9} mice (M), Alzheimer's disease	Antibiotic cocktail	Gentamicin (0.02 mg/ml), vancomycin (0.01 mg/ml), metronidazole (0.04 mg/ml), neomycin (0.01 mg/ml), ampicillin (0.02 mg/ml), kanamycin (0.06 mg/ml), colistin (120 U/ml) and cefaperazone (0.02 mg/ml) in drinking water until 6 months of age	Reduced Aβ deposition, but increased soluble Aβ levels	(Minter et al. 2016)
3xTg-AD mice (M), Alzheimer's disease	SLAB51: <i>S. thermophiles</i> , <i>B. longum</i> , <i>B. breve</i> , <i>B. infantis</i> , <i>L. acidophilus</i> , <i>L. plantarum</i> , <i>L. paracasei</i> , <i>L. delbrueckii</i> <i>subsp. bulgaricus</i> , <i>L. brevis</i>	2 × 10 ¹¹ CFU /kg in drinking water bodyweight per day	Reduced cognitive decline, brain damage, and Aβ deposition	(Bonfili et al. 2017, Bonfili et al. 2018)
Thy1-α-synuclein mice (M), Parkinson's Disease	Antibiotic cocktail	Ampicillin (1 g/L), vancomycin (0.5 g/L), neomycin (0.5 g/L), gentamycin (100 mg/L), and erythromycin (10 mg/L) in drinking water for 7 weeks	Ameliorated motor function	(Sampson et al. 2016)
Thy1-α-synuclein mice (M), Parkinson's Disease	Short-chain fatty acids	Sodium acetate (67.5mM), sodium propionate (25mM), and sodium butyrate (40mM) in drinking water for 7 weeks	Increased microglia activation and motor dysfunction	(Sampson et al. 2016)

Abbreviations: M, males; F, females; CFU, colony-forming units; LPS, lipopolysaccharide; GOS, galacto-oligosaccharide; FOS, fructo-oligosaccharide; A β , amyloid beta.

Competing interests

J.F.C. and T.G.D. received research funding from Dupont Nutrition, Cremo SA, Alkermes Inc., 4D Pharma PLC, Mead Johnson Nutrition and Nutricia Danone.

Funding

The APC Microbiome Institute is a research institute funded by Science Foundation Ireland (SFI) through the Irish Government's National Development Plan (Grant No. SFI/12/RC/2273), and SFI grant AMBROSIAC (15/JP-HDHL/3270).

References

- Aguilar-Toalá, J. E., R. Garcia-Varela, H. S. Garciac, V. Mata-Harod, A. F. González-Córdova, B. Vallejo-Cordoba and A. Hernández-Mendoza. (2018). Postbiotics: An evolving term within the functional foods field. *Trends in Food Science & Technology*, 75, 105-114. doi:10.1016/j.tifs.2018.03.009
- Aho, V. T. E., P. A. B. Pereira, S. Voutilainen, L. Paulin, E. Pekkonen, P. Auvinen and F. Scheperjans. (2019). Gut microbiota in Parkinson's disease: Temporal stability and relations to disease progression. *EBioMedicine*, 44, 691-707. doi:10.1016/j.ebiom.2019.05.064
- Ajami, B., J. L. Bennett, C. Krieger, K. M. McNagny and F. M. Rossi. (2011). Infiltrating monocytes trigger EAE progression, but do not contribute to the resident microglia pool. *Nat Neurosci*, 14(9), 1142-1149. doi:10.1038/nn.2887
- Ajami, B., J. L. Bennett, C. Krieger, W. Tetzlaff and F. M. Rossi. (2007). Local self-renewal can sustain CNS microglia maintenance and function throughout adult life. *Nat Neurosci*, 10(12), 1538-1543. doi:10.1038/nn2014
- Akbari, E., Z. Asemi, R. Daneshvar Kakhaki, F. Bahmani, E. Kouchaki, O. R. Tamtaji, G. A. Hamidi and M. Salami. (2016). Effect of Probiotic Supplementation on Cognitive Function and Metabolic Status in Alzheimer's Disease: A Randomized, Double-Blind and Controlled Trial. *Front Aging Neurosci*, 8, 256. doi:10.3389/fnagi.2016.00256
- Allen, A. P., W. Hutch, Y. E. Borre, P. J. Kennedy, A. Temko, G. Boylan, E. Murphy, J. F. Cryan, T. G. Dinan and G. Clarke. (2016). Bifidobacterium longum 1714 as a translational psychobiotic: modulation of stress, electrophysiology and neurocognition in healthy volunteers. *Transl Psychiatry*, 6(11), e939. doi:10.1038/tp.2016.191
- Andersson, H., C. Tullberg, S. Ahrne, K. Hamberg, I. Lazou Ahren, G. Molin, M. Sonesson and A. Hakansson. (2016). Oral Administration of Lactobacillus plantarum 299v Reduces Cortisol Levels in Human Saliva during Examination Induced Stress: A Randomized, Double-Blind Controlled Trial. *Int J Microbiol*, 2016, 8469018. doi:10.1155/2016/8469018
- Andonegui, G., E. L. Zelinski, C. L. Schubert, D. Knight, L. A. Craig, B. W. Winston, S. C. Spanswick, B. Petri, C. N. Jenne, J. C. Sutherland, R. Nguyen, N. Jayawardena, M. M. Kelly, C. J. Doig, R. J. Sutherland and P. Kubes. (2018). Targeting inflammatory monocytes in sepsis-associated encephalopathy and long-term cognitive impairment. *JCI Insight*, 3(9). doi:10.1172/jci.insight.99364
- Ang, Z., J. Z. Er, N. S. Tan, J. Lu, Y. C. Liou, J. Grosse and J. L. Ding. (2016). Human and mouse monocytes display distinct signalling and cytokine profiles upon stimulation with FFAR2/FFAR3 short-chain fatty acid receptor agonists. *Sci Rep*, 6, 34145. doi:10.1038/srep34145
- Arzt, E., M. Costas, S. Finkelman and V. E. Nahmod. (1991). Serotonin inhibition of tumor necrosis factor-alpha synthesis by human monocytes. *Life Sci*, 48(26), 2557-2562.
- Atashrazm, F., D. Hammond, G. Perera, C. Dobson-Stone, N. Mueller, R. Pickford, W. S. Kim, J. B. Kwok, S. J. G. Lewis, G. M. Halliday and N. Dzamko. (2018). Reduced glucocerebrosidase activity in monocytes from patients with Parkinson's disease. *Sci Rep*, 8(1), 15446. doi:10.1038/s41598-018-33921-x
- Auffray, C., D. Fogg, M. Garfa, G. Elain, O. Join-Lambert, S. Kayal, S. Sarnacki, A. Cumano, G. Lauvau and F. Geissmann. (2007). Monitoring of blood vessels and tissues by a population of monocytes with patrolling behavior. *Science*, 317(5838), 666-670. doi:10.1126/science.1142883
- Bailey, M. T., S. E. Dowd, J. D. Galley, A. R. Hufnagle, R. G. Allen and M. Lyte. (2011). Exposure to a social stressor alters the structure of the intestinal microbiota: implications for stressor-induced immunomodulation. *Brain Behav Immun*, 25(3), 397-407. doi:10.1016/j.bbi.2010.10.023
- Bailey, M. T., H. Engler, N. D. Powell, D. A. Padgett and J. F. Sheridan. (2007). Repeated social defeat increases the bactericidal activity of splenic macrophages through a Toll-like receptor-dependent pathway. *Am J Physiol Regul Integr Comp Physiol*, 293(3), R1180-1190. doi:10.1152/ajpregu.00307.2007
- Bakker, G. J. and M. Nieuwdorp. (2017). Fecal Microbiota Transplantation: Therapeutic Potential for a Multitude of Diseases beyond Clostridium difficile. *Microbiol Spectr*, 5(4). doi:10.1128/microbiolspec.BAD-0008-2017

Balmer, M. L., C. M. Schurch, Y. Saito, M. B. Geuking, H. Li, M. Cuenca, L. V. Kovtonyuk, K. D. McCoy, S. Hapfelmeier, A. F. Ochsenbein, M. G. Manz, E. Slack and A. J. Macpherson. (2014). Microbiota-derived compounds drive steady-state granulopoiesis via MyD88/TICAM signaling. *J Immunol*, 193(10), 5273-5283. doi:10.4049/jimmunol.1400762

Barichella, M., M. Severgnini, R. Cilia, E. Cassani, C. Bolliri, S. Caronni, V. Ferri, R. Canello, C. Ceccarani, S. Faierman, G. Pinelli, G. De Bellis, L. Zecca, E. Cereda, C. Consolandi and G. Pezzoli. (2019). Unraveling gut microbiota in Parkinson's disease and atypical parkinsonism. *Mov Disord*, 34(3), 396-405. doi:10.1002/mds.27581

Barrera-Bugueno, C., O. Realini, J. Escobar-Luna, R. Sotomayor-Zarate, M. Gotteland, M. Julio-Pieper and J. A. Bravo. (2017). Anxiogenic effects of a Lactobacillus, inulin and the synbiotic on healthy juvenile rats. *Neuroscience*, 359, 18-29. doi:10.1016/j.neuroscience.2017.06.064

Baufeld, C., E. O'Loughlin, N. Calcagno, C. Madore and O. Butovsky. (2018). Differential contribution of microglia and monocytes in neurodegenerative diseases. *J Neural Transm (Vienna)*, 125(5), 809-826. doi:10.1007/s00702-017-1795-7

Benakis, C., D. Brea, S. Caballero, G. Faraco, J. Moore, M. Murphy, G. Sita, G. Racchumi, L. Ling, E. G. Pamer, C. Iadecola and J. Anrather. (2016). Commensal microbiota affects ischemic stroke outcome by regulating intestinal gammadelta T cells. *Nat Med*, 22(5), 516-523. doi:10.1038/nm.4068

Berer, K., L. A. Gerdes, E. Cekanaviciute, X. Jia, L. Xiao, Z. Xia, C. Liu, L. Klotz, U. Stauter, S. E. Baranzini, T. Kumpfel, R. Hohlfeld, G. Krishnamoorthy and H. Wekerle. (2017). Gut microbiota from multiple sclerosis patients enables spontaneous autoimmune encephalomyelitis in mice. *Proc Natl Acad Sci U S A*, 114(40), 10719-10724. doi:10.1073/pnas.1711233114

Bettcher, B. M., J. Neuhaus, M. J. Wynn, F. M. Elahi, K. B. Casaleto, R. Saloner, R. Fitch, A. Karydas and J. H. Kramer. (2019). Increases in a Pro-inflammatory Chemokine, MCP-1, Are Related to Decreases in Memory Over Time. *Front Aging Neurosci*, 11, 25. doi:10.3389/fnagi.2019.00025

Bharwani, A., M. F. Mian, M. G. Surette, J. Bienenstock and P. Forsythe. (2017). Oral treatment with Lactobacillus rhamnosus attenuates behavioural deficits and immune changes in chronic social stress. *BMC Med*, 15(1), 7. doi:10.1186/s12916-016-0771-7

Bhatnagar, S., C. Vining, V. Iyer and V. Kinni. (2006). Changes in hypothalamic-pituitary-adrenal function, body temperature, body weight and food intake with repeated social stress exposure in rats. *J Neuroendocrinol*, 18(1), 13-24. doi:10.1111/j.1365-2826.2005.01375.x

Biagi, E., L. Nylund, M. Candela, R. Ostan, L. Bucci, E. Pini, J. Nikkila, D. Monti, R. Satokari, C. Franceschi, P. Brigidi and W. De Vos. (2010). Through ageing, and beyond: gut microbiota and inflammatory status in seniors and centenarians. *PLoS One*, 5(5), e10667. doi:10.1371/journal.pone.0010667

Bindels, L. B., N. M. Delzenne, P. D. Cani and J. Walter. (2015). Towards a more comprehensive concept for prebiotics. *Nat Rev Gastroenterol Hepatol*, 12(5), 303-310. doi:10.1038/nrgastro.2015.47

Bloemen, J. G., K. Venema, M. C. van de Poll, S. W. Olde Damink, W. A. Buurman and C. H. Dejong. (2009). Short chain fatty acids exchange across the gut and liver in humans measured at surgery. *Clin Nutr*, 28(6), 657-661. doi:10.1016/j.clnu.2009.05.011

Boehme, M., M. van de Wouw, T. F. S. Bastiaansen, L. Olavarria-Ramirez, K. Lyons, F. Fouhy, A. V. Golubeva, G. M. Moloney, C. Minuto, K. V. Sandhu, K. A. Scott, G. Clarke, C. Stanton, T. G. Dinan, H. Schellekens and J. F. Cryan. (2019). Mid-life microbiota crises: middle age is associated with pervasive neuroimmune alterations that are reversed by targeting the gut microbiome. *Mol Psychiatry*. doi:10.1038/s41380-019-0425-1

Bonaz, B., T. Bazin and S. Pellissier. (2018). The Vagus Nerve at the Interface of the Microbiota-Gut-Brain Axis. *Front Neurosci*, 12, 49. doi:10.3389/fnins.2018.00049

Bonfili, L., V. Cecarini, S. Berardi, S. Scarpona, J. S. Suchodolski, C. Nasuti, D. Fiorini, M. C. Boarelli, G. Rossi and A. M. Eleuteri. (2017). Microbiota modulation counteracts Alzheimer's disease progression influencing neuronal proteolysis and gut hormones plasma levels. *Sci Rep*, 7(1), 2426. doi:10.1038/s41598-017-02587-2

Bonfili, L., V. Cecarini, M. Cuccioloni, M. Angeletti, S. Berardi, S. Scarpona, G. Rossi and A. M. Eleuteri. (2018). SLAB51 Probiotic Formulation Activates SIRT1 Pathway Promoting Antioxidant and

Neuroprotective Effects in an AD Mouse Model. *Mol Neurobiol*, 55(10), 7987-8000. doi:10.1007/s12035-018-0973-4

Bordon, Y. (2014). Macrophages: innate memory training. *Nat Rev Immunol*, 14(11), 713. doi:10.1038/nri3759

Brattsand, R. and M. Linden. (1996). Cytokine modulation by glucocorticoids: mechanisms and actions in cellular studies. *Aliment Pharmacol Ther*, 10 Suppl 2, 81-90; discussion 91-82.

Bravo, J. A., P. Forsythe, M. V. Chew, E. Escaravage, H. M. Savignac, T. G. Dinan, J. Bienenstock and J. F. Cryan. (2011). Ingestion of Lactobacillus strain regulates emotional behavior and central GABA receptor expression in a mouse via the vagus nerve. *Proc Natl Acad Sci U S A*, 108(38), 16050-16055. doi:10.1073/pnas.1102999108

Brazaitis, M., N. Eimantas, L. Daniuseviciute, D. Mickeviciene, R. Steponaviciute and A. Skurvydas. (2014). Two strategies for response to 14 degrees C cold-water immersion: is there a difference in the response of motor, cognitive, immune and stress markers? *PLoS One*, 9(9), e109020. doi:10.1371/journal.pone.0109020

Brenner, L. A., K. A. Stearns-Yoder, A. S. Hoffberg, M. E. Penzenik, A. J. Starosta, T. D. Hernandez, D. A. Hadidi and C. A. Lowry. (2017). Growing literature but limited evidence: A systematic review regarding prebiotic and probiotic interventions for those with traumatic brain injury and/or posttraumatic stress disorder. *Brain Behav Immun*, 65, 57-67. doi:10.1016/j.bbi.2017.06.003

Bruunsgaard, H., K. Andersen-Ranberg, B. Jeune, A. N. Pedersen, P. Skinhoj and B. K. Pedersen. (1999). A high plasma concentration of TNF-alpha is associated with dementia in centenarians. *J Gerontol A Biol Sci Med Sci*, 54(7), M357-364.

Burokas, A., S. Arboleya, R. D. Moloney, V. L. Peterson, K. Murphy, G. Clarke, C. Stanton, T. G. Dinan and J. F. Cryan. (2017). Targeting the Microbiota-Gut-Brain Axis: Prebiotics Have Anxiolytic and Antidepressant-like Effects and Reverse the Impact of Chronic Stress in Mice. *Biol Psychiatry*, 82(7), 472-487. doi:10.1016/j.biopsych.2016.12.031

Butovsky, O., M. P. Jedrychowski, C. S. Moore, R. Cialic, A. J. Lanser, G. Gabriely, T. Koeglspenger, B. Dake, P. M. Wu, C. E. Doykan, Z. Fanek, L. Liu, Z. Chen, J. D. Rothstein, R. M. Ransohoff, S. P. Gygi, J. P. Antel and H. L. Weiner. (2014). Identification of a unique TGF-beta-dependent molecular and functional signature in microglia. *Nat Neurosci*, 17(1), 131-143. doi:10.1038/nn.3599

Campbell, C. L., R. Yu, F. Li, Q. Zhou, D. Chen, C. Qi, Y. Yin and J. Sun. (2019). Modulation of fat metabolism and gut microbiota by resveratrol on high-fat diet-induced obese mice. *Diabetes Metab Syndr Obes*, 12, 97-107. doi:10.2147/DMSO.S192228

Canivet, C., O. Uyar, C. Rheaume, J. Piret and G. Boivin. (2019). The recruitment of peripheral blood leukocytes to the brain is delayed in susceptible BALB/c compared to resistant C57BL/6 mice during herpes simplex virus encephalitis. *J Neurovirol*. doi:10.1007/s13365-019-00730-5

Cavaillon, J. M. (2018). Exotoxins and endotoxins: Inducers of inflammatory cytokines. *Toxicon*, 149, 45-53. doi:10.1016/j.toxicon.2017.10.016

Cekanaviciute, E., B. B. Yoo, T. F. Runia, J. W. Debelius, S. Singh, C. A. Nelson, R. Kanner, Y. Bencosme, Y. K. Lee, S. L. Hauser, E. Crabtree-Hartman, I. K. Sand, M. Gacias, Y. Zhu, P. Casaccia, B. A. C. Cree, R. Knight, S. K. Mazmanian and S. E. Baranzini. (2017). Gut bacteria from multiple sclerosis patients modulate human T cells and exacerbate symptoms in mouse models. *Proc Natl Acad Sci U S A*, 114(40), 10713-10718. doi:10.1073/pnas.1711235114

Chen, J., N. Chia, K. R. Kalari, J. Z. Yao, M. Novotna, M. M. Paz Soldan, D. H. Luckey, E. V. Marietta, P. R. Jeraldo, X. Chen, B. G. Weinshenker, M. Rodriguez, O. H. Kantarci, H. Nelson, J. A. Murray and A. K. Mangalam. (2016). Multiple sclerosis patients have a distinct gut microbiota compared to healthy controls. *Sci Rep*, 6, 28484. doi:10.1038/srep28484

Chen, M. L., L. Yi, Y. Zhang, X. Zhou, L. Ran, J. Yang, J. D. Zhu, Q. Y. Zhang and M. T. Mi. (2016). Resveratrol Attenuates Trimethylamine-N-Oxide (TMAO)-Induced Atherosclerosis by Regulating TMAO Synthesis and Bile Acid Metabolism via Remodeling of the Gut Microbiota. *MBio*, 7(2), e02210-02215. doi:10.1128/mBio.02210-15

Cheng, S. C., J. Quintin, R. A. Cramer, K. M. Shepardson, S. Saeed, V. Kumar, E. J. Giamarellos-Bourboulis, J. H. Martens, N. A. Rao, A. Aghajani-Refah, G. R. Manjeri, Y. Li, D. C. Ifrim, R. J. Arts, B. M. van der Veer, P. M. Deen, C. Logie, L. A. O'Neill, P. Willems, F. L. van de Veerdonk, J. W. van der Meer, A. Ng, L. A. Joosten, C. Wijmenga, H. G. Stunnenberg, R. J. Xavier and M. G. Netea. (2014). mTOR- and HIF-1 α -mediated aerobic glycolysis as metabolic basis for trained immunity. *Science*, 345(6204), 1250684. doi:10.1126/science.1250684

Chou, A., K. Krukowski, J. M. Morganti, L. K. Riparip and S. Rosi. (2018). Persistent Infiltration and Impaired Response of Peripherally-Derived Monocytes after Traumatic Brain Injury in the Aged Brain. *Int J Mol Sci*, 19(6). doi:10.3390/ijms19061616

Chung, Y. C., H. M. Jin, Y. Cui, D. S. Kim, J. M. Jung, J. I. Park, E. S. Jung, E. K. Choi and S. W. Chae. (2014). Fermented milk of *Lactobacillus helveticus* IDCC3801 improves cognitive functioning during cognitive fatigue tests in healthy older adults. *Journal of Functional Foods*, 10, 465-474. doi:10.1016/j.jff.2014.07.007

Cignarella, F., C. Cantoni, L. Ghezzi, A. Salter, Y. Dorsett, L. Chen, D. Phillips, G. M. Weinstock, L. Fontana, A. H. Cross, Y. Zhou and L. Piccio. (2018). Intermittent Fasting Confers Protection in CNS Autoimmunity by Altering the Gut Microbiota. *Cell Metab*, 27(6), 1222-1235 e1226. doi:10.1016/j.cmet.2018.05.006

Claesson, M. J., I. B. Jeffery, S. Conde, S. E. Power, E. M. O'Connor, S. Cusack, H. M. Harris, M. Coakley, B. Lakshminarayanan, O. O'Sullivan, G. F. Fitzgerald, J. Deane, M. O'Connor, N. Harnedy, K. O'Connor, D. O'Mahony, D. van Sinderen, M. Wallace, L. Brennan, C. Stanton, J. R. Marchesi, A. P. Fitzgerald, F. Shanahan, C. Hill, R. P. Ross and P. W. O'Toole. (2012). Gut microbiota composition correlates with diet and health in the elderly. *Nature*, 488(7410), 178-184. doi:10.1038/nature11319

Clarke, G., S. Grenham, P. Scully, P. Fitzgerald, R. D. Moloney, F. Shanahan, T. G. Dinan and J. F. Cryan. (2013). The microbiome-gut-brain axis during early life regulates the hippocampal serotonergic system in a sex-dependent manner. *Mol Psychiatry*, 18(6), 666-673. doi:10.1038/mp.2012.77

Cleophas, M. C. P., J. M. Ratter, S. Bekkering, J. Quintin, K. Schraa, E. S. Stroes, M. G. Netea and L. A. B. Joosten. (2019). Effects of oral butyrate supplementation on inflammatory potential of circulating peripheral blood mononuclear cells in healthy and obese males. *Sci Rep*, 9(1), 775. doi:10.1038/s41598-018-37246-7

Cloez-Tayarani, I., A. F. Petit-Bertron, H. D. Venters and J. M. Cavaillon. (2003). Differential effect of serotonin on cytokine production in lipopolysaccharide-stimulated human peripheral blood mononuclear cells: involvement of 5-hydroxytryptamine_{2A} receptors. *Int Immunol*, 15(2), 233-240.

Collins, S. M., M. Surette and P. Bercik. (2012). The interplay between the intestinal microbiota and the brain. *Nat Rev Microbiol*, 10(11), 735-742. doi:10.1038/nrmicro2876

Columba-Cabezas, S., B. Serafini, E. Ambrosini, M. Sanchez, G. Penna, L. Adorini and F. Aloisi. (2002). Induction of macrophage-derived chemokine/CCL22 expression in experimental autoimmune encephalomyelitis and cultured microglia: implications for disease regulation. *J Neuroimmunol*, 130(1-2), 10-21.

Cooper, M. A. and K. L. Huhman. (2010). Blocking corticotropin-releasing factor-2 receptors, but not corticotropin-releasing factor-1 receptors or glucocorticoid feedback, disrupts the development of conditioned defeat. *Physiol Behav*, 101(4), 527-532. doi:10.1016/j.physbeh.2010.08.003

Costantini, L., R. Molinari, B. Farinon and N. Merendino. (2017). Impact of Omega-3 Fatty Acids on the Gut Microbiota. *Int J Mol Sci*, 18(12). doi:10.3390/ijms18122645

Coutinho, A. E. and K. E. Chapman. (2011). The anti-inflammatory and immunosuppressive effects of glucocorticoids, recent developments and mechanistic insights. *Mol Cell Endocrinol*, 335(1), 2-13. doi:10.1016/j.mce.2010.04.005

Crapser, J., R. Ritzel, R. Verma, V. R. Venna, F. Liu, A. Chauhan, E. Koellhoffer, A. Patel, A. Ricker, K. Maas, J. Graf and L. D. McCullough. (2016). Ischemic stroke induces gut permeability and enhances bacterial translocation leading to sepsis in aged mice. *Aging (Albany NY)*, 8(5), 1049-1063. doi:10.18632/aging.100952

Cronk, J. C., A. J. Filiano, A. Louveau, I. Marin, R. Marsh, E. Ji, D. H. Goldman, I. Smirnov, N. Geraci, S. Acton, C. C. Overall and J. Kipnis. (2018). Peripherally derived macrophages can engraft the brain independent of irradiation and maintain an identity distinct from microglia. *J Exp Med*, 215(6), 1627-1647. doi:10.1084/jem.20180247

Cryan, J. F. and T. G. Dinan. (2012). Mind-altering microorganisms: the impact of the gut microbiota on brain and behaviour. *Nat Rev Neurosci*, 13(10), 701-712. doi:10.1038/nrn3346

Cryan, J. F., K. J. O'Riordan, C. S. M. Cowan, K. V. Sandhu, T. F. S. Bastiaanssen, M. Boehme, M. G. Codagnone, S. Cusotto, C. Fulling, A. V. Golubeva, K. E. Guzzetta, M. Jagger, C. M. Long-Smith, J. M. Lyte, J. A. Martin, A. Molinero-Perez, G. Moloney, E. Morelli, E. Morillas, R. O'Connor, J. Pereira, V. L. Peterson, K. Rea, N. L. Ritz, E. Sherwin, S. Spichak, E. M. Teichman, M. Van de Wouw, A. P. Ventura-Silva, S. E. Wallace-Fitzsimons, N. P. Hyland, G. Clarke and T. G. Dinan. (2019). The Microbiota-Gut-Brain Axis. *Accepted in Physiological Reviews*.

Cummings, J. H., E. W. Pomare, W. J. Branch, C. P. Naylor and G. T. Macfarlane. (1987). Short chain fatty acids in human large intestine, portal, hepatic and venous blood. *Gut*, 28(10), 1221-1227.

D'Mello, C., W. Almishri, H. Liu and M. G. Swain. (2017). Interactions Between Platelets and Inflammatory Monocytes Affect Sickness Behavior in Mice With Liver Inflammation. *Gastroenterology*, 153(5), 1416-1428 e1412. doi:10.1053/j.gastro.2017.08.011

D'Mello, C., T. Le and M. G. Swain. (2009). Cerebral microglia recruit monocytes into the brain in response to tumor necrosis factor- α signaling during peripheral organ inflammation. *J Neurosci*, 29(7), 2089-2102. doi:10.1523/JNEUROSCI.3567-08.2009

D'Mello, C., K. Riaz, T. Le, K. M. Stevens, A. Wang, D. M. McKay, Q. J. Pittman and M. G. Swain. (2013). P-selectin-mediated monocyte-cerebral endothelium adhesive interactions link peripheral organ inflammation to sickness behaviors. *J Neurosci*, 33(37), 14878-14888. doi:10.1523/JNEUROSCI.1329-13.2013

D'Mello, C., N. Ronaghan, R. Zaheer, M. Dicay, T. Le, W. K. MacNaughton, M. G. Surette and M. G. Swain. (2015). Probiotics Improve Inflammation-Associated Sickness Behavior by Altering Communication between the Peripheral Immune System and the Brain. *J Neurosci*, 35(30), 10821-10830. doi:10.1523/JNEUROSCI.0575-15.2015

Dalmasso, M., C. Hill and R. P. Ross. (2014). Exploiting gut bacteriophages for human health. *Trends Microbiol*, 22(7), 399-405. doi:10.1016/j.tim.2014.02.010

Dantzer, R., C. J. Heijnen, A. Kavelaars, S. Laye and L. Capuron. (2014). The neuroimmune basis of fatigue. *Trends Neurosci*, 37(1), 39-46. doi:10.1016/j.tins.2013.10.003

Dantzer, R., J. C. O'Connor, G. G. Freund, R. W. Johnson and K. W. Kelley. (2008). From inflammation to sickness and depression: when the immune system subjugates the brain. *Nat Rev Neurosci*, 9(1), 46-56. doi:10.1038/nrn2297

de las Casas-Engel, M., A. Dominguez-Soto, E. Sierra-Filardi, R. Bragado, C. Nieto, A. Puig-Kroger, R. Samaniego, M. Loza, M. T. Corcuera, F. Gomez-Aguado, M. Bustos, P. Sanchez-Mateos and A. L. Corbi. (2013). Serotonin skews human macrophage polarization through HTR2B and HTR7. *J Immunol*, 190(5), 2301-2310. doi:10.4049/jimmunol.1201133

Derrien, M. and J. E. van Hylckama Vlieg. (2015). Fate, activity, and impact of ingested bacteria within the human gut microbiota. *Trends Microbiol*, 23(6), 354-366. doi:10.1016/j.tim.2015.03.002

Dhabhar, F. S., W. B. Malarkey, E. Neri and B. S. McEwen. (2012). Stress-induced redistribution of immune cells--from barracks to boulevards to battlefields: a tale of three hormones--Curt Richter Award winner. *Psychoneuroendocrinology*, 37(9), 1345-1368. doi:10.1016/j.psyneuen.2012.05.008

Dhabhar, F. S. and B. S. McEwen. (1996). Stress-induced enhancement of antigen-specific cell-mediated immunity. *J Immunol*, 156(7), 2608-2615.

Dhabhar, F. S., A. H. Miller, B. S. McEwen and R. L. Spencer. (1995). Effects of stress on immune cell distribution. Dynamics and hormonal mechanisms. *J Immunol*, 154(10), 5511-5527.

Dhabhar, F. S., A. H. Miller, M. Stein, B. S. McEwen and R. L. Spencer. (1994). Diurnal and acute stress-induced changes in distribution of peripheral blood leukocyte subpopulations. *Brain Behav Immun*, 8(1), 66-79. doi:10.1006/brbi.1994.1006

Dhaliwal, J., D. P. Singh, S. Singh, A. K. Pinnaka, R. K. Boparai, M. Bishnoi, K. K. Kondepudi and K. Chopra. (2018). Lactobacillus plantarum MTCC 9510 supplementation protects from chronic unpredictable and sleep deprivation-induced behaviour, biochemical and selected gut microbial aberrations in mice. *J Appl Microbiol*, 125(1), 257-269. doi:10.1111/jam.13765

Dinan, T. G., C. Stanton and J. F. Cryan. (2013). Psychobiotics: a novel class of psychotropic. *Biol Psychiatry*, 74(10), 720-726. doi:10.1016/j.biopsych.2013.05.001

Diop, L., S. Guillou and H. Durand. (2008). Probiotic food supplement reduces stress-induced gastrointestinal symptoms in volunteers: a double-blind, placebo-controlled, randomized trial. *Nutr Res*, 28(1), 1-5. doi:10.1016/j.nutres.2007.10.001

Drevets, D. A., M. J. Dillon, J. S. Schawang, N. Van Rooijen, J. Ehrchen, C. Sunderkotter and P. J. Leenen. (2004). The Ly-6Chigh monocyte subpopulation transports *Listeria monocytogenes* into the brain during systemic infection of mice. *J Immunol*, 172(7), 4418-4424.

Dunphy-Doherty, F., S. M. O'Mahony, V. L. Peterson, O. O'Sullivan, F. Crispie, P. D. Cotter, P. Wigmore, M. V. King, J. F. Cryan and K. C. F. Fone. (2018). Post-weaning social isolation of rats leads to long-term disruption of the gut microbiota-immune-brain axis. *Brain Behav Immun*, 68, 261-273. doi:10.1016/j.bbi.2017.10.024

Durk, T., E. Panther, T. Muller, S. Sorichter, D. Ferrari, C. Pizzirani, F. Di Virgilio, D. Myrtek, J. Norgauer and M. Idzko. (2005). 5-Hydroxytryptamine modulates cytokine and chemokine production in LPS-primed human monocytes via stimulation of different 5-HTR subtypes. *Int Immunol*, 17(5), 599-606. doi:10.1093/intimm/dxh242

Ehrchen, J., L. Steinmuller, K. Barczyk, K. Tenbrock, W. Nacken, M. Eisenacher, U. Nordhues, C. Sorg, C. Sunderkotter and J. Roth. (2007). Glucocorticoids induce differentiation of a specifically activated, anti-inflammatory subtype of human monocytes. *Blood*, 109(3), 1265-1274. doi:10.1182/blood-2006-02-001115

El Khoury, J., M. Toft, S. E. Hickman, T. K. Means, K. Terada, C. Geula and A. D. Luster. (2007). *Ccr2* deficiency impairs microglial accumulation and accelerates progression of Alzheimer-like disease. *Nat Med*, 13(4), 432-438. doi:10.1038/nm1555

Engler, H., M. T. Bailey, A. Engler and J. F. Sheridan. (2004). Effects of repeated social stress on leukocyte distribution in bone marrow, peripheral blood and spleen. *J Neuroimmunol*, 148(1-2), 106-115. doi:10.1016/j.jneuroim.2003.11.011

Engler, H., M. T. Bailey, A. Engler, L. M. Stiner-Jones, N. Quan and J. F. Sheridan. (2008). Interleukin-1 receptor type 1-deficient mice fail to develop social stress-associated glucocorticoid resistance in the spleen. *Psychoneuroendocrinology*, 33(1), 108-117. doi:10.1016/j.psyneuen.2007.10.007

Engler, H., A. Engler, M. T. Bailey and J. F. Sheridan. (2005). Tissue-specific alterations in the glucocorticoid sensitivity of immune cells following repeated social defeat in mice. *J Neuroimmunol*, 163(1-2), 110-119. doi:10.1016/j.jneuroim.2005.03.002

Erny, D., A. L. Hrabé de Angelis, D. Jaitin, P. Wieghofer, O. Staszewski, E. David, H. Keren-Shaul, T. Mhlakoiv, K. Jakobshagen, T. Buch, V. Schwierzeck, O. Utermohlen, E. Chun, W. S. Garrett, K. D. McCoy, A. Diefenbach, P. Staeheli, B. Stecher, I. Amit and M. Prinz. (2015). Host microbiota constantly control maturation and function of microglia in the CNS. *Nat Neurosci*, 18(7), 965-977. doi:10.1038/nn.4030

Falcão de Arruda, I. S. and J. E. de Aguilar-Nascimento. (2004). Benefits of early enteral nutrition with glutamine and probiotics in brain injury patients. *Clin Sci (Lond)*, 106(3), 287-292. doi:10.1042/CS20030251

Fauci, A. S. and D. C. Dale. (1974). The effect of in vivo hydrocortisone on subpopulations of human lymphocytes. *J Clin Invest*, 53(1), 240-246. doi:10.1172/JCI107544

Fife, B. T., G. B. Huffnagle, W. A. Kuziel and W. J. Karpus. (2000). CC chemokine receptor 2 is critical for induction of experimental autoimmune encephalomyelitis. *J Exp Med*, 192(6), 899-905.

Finger, B. C., T. G. Dinan and J. F. Cryan. (2011). High-fat diet selectively protects against the effects of chronic social stress in the mouse. *Neuroscience*, 192, 351-360. doi:10.1016/j.neuroscience.2011.06.072

Fischer, H. J., T. L. K. Finck, H. L. Pellkofer, H. M. Reichardt and F. Luhder. (2019). Glucocorticoid Therapy of Multiple Sclerosis Patients Induces Anti-inflammatory Polarization and Increased Chemotaxis of Monocytes. *Front Immunol*, 10, 1200. doi:10.3389/fimmu.2019.01200

Foertsch, S., A. M. Fuchsl, S. D. Faller, H. Holzer, D. Langgartner, J. Messmann, G. Strauss and S. O. Reber. (2017). Splenic glucocorticoid resistance following psychosocial stress requires physical injury. *Sci Rep*, 7(1), 15730. doi:10.1038/s41598-017-15897-2

Franceschi, C., M. Bonafe, S. Valensin, F. Olivieri, M. De Luca, E. Ottaviani and G. De Benedictis. (2000). Inflamm-aging. An evolutionary perspective on immunosenescence. *Ann N Y Acad Sci*, 908, 244-254.

Frank, M. G., Z. D. Miguel, L. R. Watkins and S. F. Maier. (2010). Prior exposure to glucocorticoids sensitizes the neuroinflammatory and peripheral inflammatory responses to *E. coli* lipopolysaccharide. *Brain Behav Immun*, 24(1), 19-30. doi:10.1016/j.bbi.2009.07.008

Fransen, F., A. A. van Beek, T. Borghuis, S. E. Aidy, F. Hugenholtz, C. van der Gaast-de Jongh, H. F. J. Savelkoul, M. I. De Jonge, M. V. Boekschoten, H. Smidt, M. M. Faas and P. de Vos. (2017). Aged Gut Microbiota Contributes to Systemic Inflammation after Transfer to Germ-Free Mice. *Front Immunol*, 8, 1385. doi:10.3389/fimmu.2017.01385

Frasca, D. and B. B. Blomberg. (2016). Inflammation decreases adaptive and innate immune responses in mice and humans. *Biogerontology*, 17(1), 7-19. doi:10.1007/s10522-015-9578-8

Fuchsl, A. M., D. Langgartner and S. O. Reber. (2013). Mechanisms underlying the increased plasma ACTH levels in chronic psychosocially stressed male mice. *PLoS One*, 8(12), e84161. doi:10.1371/journal.pone.0084161

Fukui, H., T. Oshima, Y. Tanaka, Y. Oikawa, Y. Makizaki, H. Ohno, T. Tomita, J. Watari and H. Miwa. (2018). Effect of probiotic *Bifidobacterium bifidum* G9-1 on the relationship between gut microbiota profile and stress sensitivity in maternally separated rats. *Sci Rep*, 8(1), 12384. doi:10.1038/s41598-018-30943-3

Fulling, C., T. G. Dinan and J. F. Cryan. (2019). Gut Microbe to Brain Signaling: What Happens in Vagus. *Neuron*, 101(6), 998-1002. doi:10.1016/j.neuron.2019.02.008

Galley, J. D., A. R. Mackos, V. A. Varaljay and M. T. Bailey. (2017). Stressor exposure has prolonged effects on colonic microbial community structure in *Citrobacter rodentium*-challenged mice. *Sci Rep*, 7, 45012. doi:10.1038/srep45012

Galley, J. D., M. C. Nelson, Z. Yu, S. E. Dowd, J. Walter, P. S. Kumar, M. Lyte and M. T. Bailey. (2014). Exposure to a social stressor disrupts the community structure of the colonic mucosa-associated microbiota. *BMC Microbiol*, 14, 189. doi:10.1186/1471-2180-14-189

Garcia-Bonilla, L., D. Brea, C. Benakis, D. A. Lane, M. Murphy, J. Moore, G. Racchumi, X. Jiang, C. Iadecola and J. Anrather. (2018). Endogenous Protection from Ischemic Brain Injury by Preconditioned Monocytes. *J Neurosci*, 38(30), 6722-6736. doi:10.1523/JNEUROSCI.0324-18.2018

Garre, J. M., H. M. Silva, J. J. Lafaille and G. Yang. (2017). CX3CR1(+) monocytes modulate learning and learning-dependent dendritic spine remodeling via TNF- α . *Nat Med*, 23(6), 714-722. doi:10.1038/nm.4340

Ge, R., D. Tornero, M. Hirota, E. Monni, C. Laterza, O. Lindvall and Z. Kokaia. (2017). Choroid plexus-cerebrospinal fluid route for monocyte-derived macrophages after stroke. *J Neuroinflammation*, 14(1), 153. doi:10.1186/s12974-017-0909-3

Geissmann, F., S. Jung and D. R. Littman. (2003). Blood monocytes consist of two principal subsets with distinct migratory properties. *Immunity*, 19(1), 71-82.

Getts, D. R., R. L. Terry, M. T. Getts, M. Muller, S. Rana, B. Shrestha, J. Radford, N. Van Rooijen, I. L. Campbell and N. J. King. (2008). Ly6c+ "inflammatory monocytes" are microglial precursors recruited in a pathogenic manner in West Nile virus encephalitis. *J Exp Med*, 205(10), 2319-2337. doi:10.1084/jem.20080421

Gibson, G. R., R. Hutkins, M. E. Sanders, S. L. Prescott, R. A. Reimer, S. J. Salminen, K. Scott, C. Stanton, K. S. Swanson, P. D. Cani, K. Verbeke and G. Reid. (2017). Expert consensus document: The International Scientific Association for Probiotics and Prebiotics (ISAPP) consensus statement on the

definition and scope of prebiotics. *Nat Rev Gastroenterol Hepatol*, 14(8), 491-502. doi:10.1038/nrgastro.2017.75

Ginhoux, F., M. Greter, M. Leboeuf, S. Nandi, P. See, S. Gokhan, M. F. Mehler, S. J. Conway, L. G. Ng, E. R. Stanley, I. M. Samokhvalov and M. Merad. (2010). Fate mapping analysis reveals that adult microglia derive from primitive macrophages. *Science*, 330(6005), 841-845. doi:10.1126/science.1194637

Ginhoux, F. and S. Jung. (2014). Monocytes and macrophages: developmental pathways and tissue homeostasis. *Nat Rev Immunol*, 14(6), 392-404. doi:10.1038/nri3671

Guarda, G., M. Braun, F. Staehli, A. Tardivel, C. Mattmann, I. Forster, M. Farlik, T. Decker, R. A. Du Pasquier, P. Romero and J. Tschopp. (2011). Type I interferon inhibits interleukin-1 production and inflammasome activation. *Immunity*, 34(2), 213-223. doi:10.1016/j.immuni.2011.02.006

Guilliams, M., A. Mildner and S. Yona. (2018). Developmental and Functional Heterogeneity of Monocytes. *Immunity*, 49(4), 595-613. doi:10.1016/j.immuni.2018.10.005

Guo, H., Z. Zhao, R. Zhang, P. Chen, X. Zhang, F. Cheng and X. Gou. (2019). Monocytes in the Peripheral Clearance of Amyloid-beta and Alzheimer's Disease. *J Alzheimers Dis*, 68(4), 1391-1400. doi:10.3233/JAD-181177

Gururajan, A., M. van de Wouw, M. Boehme, T. Becker, R. O'Connor, T. F. S. Bastiaanssen, G. M. Moloney, J. M. Lyte, A. Paula Ventura Silva, B. Merckx, T. G. Dinan and J. F. Cryan. (2019). Resilience to Chronic Stress Is Associated with Specific Neurobiological, Neuroendocrine and Immune Responses. *Brain Behav Immun*. doi:10.1016/j.bbi.2019.05.004

Gyoneva, S., D. Kim, A. Katsumoto, O. N. Kokiko-Cochran, B. T. Lamb and R. M. Ransohoff. (2015). Ccr2 deletion dissociates cavity size and tau pathology after mild traumatic brain injury. *J Neuroinflammation*, 12, 228. doi:10.1186/s12974-015-0443-0

Hanke, M. L., N. D. Powell, L. M. Stiner, M. T. Bailey and J. F. Sheridan. (2012). Beta adrenergic blockade decreases the immunomodulatory effects of social disruption stress. *Brain Behav Immun*, 26(7), 1150-1159. doi:10.1016/j.bbi.2012.07.011

Hansel, A., S. Hong, R. J. Camara and R. von Kanel. (2010). Inflammation as a psychophysiological biomarker in chronic psychosocial stress. *Neurosci Biobehav Rev*, 35(1), 115-121. doi:10.1016/j.neubiorev.2009.12.012

Harach, T., N. Marungruang, N. Duthilleul, V. Cheatham, K. D. McCoy, G. Frisoni, J. J. Neher, F. Fak, M. Jucker, T. Lasser and T. Bolmont. (2017). Reduction of Abeta amyloid pathology in APPS1 transgenic mice in the absence of gut microbiota. *Sci Rep*, 7, 41802. doi:10.1038/srep41802

Harden, L. M., S. Kent, Q. J. Pittman and J. Roth. (2015). Fever and sickness behavior: Friend or foe? *Brain Behav Immun*, 50, 322-333. doi:10.1016/j.bbi.2015.07.012

Harms, A. S., A. D. Thome, Z. Yan, A. M. Schonhoff, G. P. Williams, X. Li, Y. Liu, H. Qin, E. N. Benveniste and D. G. Standaert. (2018). Peripheral monocyte entry is required for alpha-Synuclein induced inflammation and Neurodegeneration in a model of Parkinson disease. *Exp Neurol*, 300, 179-187. doi:10.1016/j.expneurol.2017.11.010

Hasegawa, S., S. Goto, H. Tsuji, T. Okuno, T. Asahara, K. Nomoto, A. Shibata, Y. Fujisawa, T. Minato, A. Okamoto, K. Ohno and M. Hirayama. (2015). Intestinal Dysbiosis and Lowered Serum Lipopolysaccharide-Binding Protein in Parkinson's Disease. *PLoS One*, 10(11), e0142164. doi:10.1371/journal.pone.0142164

Hashimoto, D., A. Chow, C. Noizat, P. Teo, M. B. Beasley, M. Leboeuf, C. D. Becker, P. See, J. Price, D. Lucas, M. Greter, A. Mortha, S. W. Boyer, E. C. Forsberg, M. Tanaka, N. van Rooijen, A. Garcia-Sastre, E. R. Stanley, F. Ginhoux, P. S. Frenette and M. Merad. (2013). Tissue-resident macrophages self-maintain locally throughout adult life with minimal contribution from circulating monocytes. *Immunity*, 38(4), 792-804. doi:10.1016/j.immuni.2013.04.004

He, B., T. K. Hoang, X. Tian, C. M. Taylor, E. Blanchard, M. Luo, M. B. Bhattacharjee, J. Freeborn, S. Park, J. Couturier, J. W. Lindsey, D. Q. Tran, J. M. Rhoads and Y. Liu. (2019). Lactobacillus reuteri Reduces the Severity of Experimental Autoimmune Encephalomyelitis in Mice by Modulating Gut Microbiota. *Front Immunol*, 10, 385. doi:10.3389/fimmu.2019.00385

Hearps, A. C., G. E. Martin, T. A. Angelovich, W. J. Cheng, A. Maisa, A. L. Landay, A. Jaworowski and S. M. Crowe. (2012). Aging is associated with chronic innate immune activation and dysregulation of monocyte phenotype and function. *Aging Cell*, 11(5), 867-875. doi:10.1111/j.1474-9726.2012.00851.x

Heidt, T., H. B. Sager, G. Courties, P. Dutta, Y. Iwamoto, A. Zaltsman, C. von Zur Muhlen, C. Bode, G. L. Fricchione, J. Denninger, C. P. Lin, C. Vinegoni, P. Libby, F. K. Swirski, R. Weissleder and M. Nahrendorf. (2014). Chronic variable stress activates hematopoietic stem cells. *Nat Med*, 20(7), 754-758. doi:10.1038/nm.3589

Hench, P. S., C. H. Slocumb, H. F. Polley and E. C. Kendal. (1950). Effect of cortisone and pituitary adrenocorticotrophic hormone (ACTH) on rheumatic diseases. *J Am Med Assoc*, 144(16), 1327-1335.

Herisson, F., V. Frodermann, G. Courties, D. Rohde, Y. Sun, K. Vandoorne, G. R. Wojtkiewicz, G. S. Masson, C. Vinegoni, J. Kim, D. E. Kim, R. Weissleder, F. K. Swirski, M. A. Moskowitz and M. Nahrendorf. (2018). Direct vascular channels connect skull bone marrow and the brain surface enabling myeloid cell migration. *Nat Neurosci*, 21(9), 1209-1217. doi:10.1038/s41593-018-0213-2

Herr, N., C. Bode and D. Duerschmied. (2017). The Effects of Serotonin in Immune Cells. *Front Cardiovasc Med*, 4, 48. doi:10.3389/fcvm.2017.00048

Hickman, S. E., N. D. Kingery, T. K. Ohsumi, M. L. Borowsky, L. C. Wang, T. K. Means and J. El Khoury. (2013). The microglial sensome revealed by direct RNA sequencing. *Nat Neurosci*, 16(12), 1896-1905. doi:10.1038/nn.3554

Hill, C., F. Guarner, G. Reid, G. R. Gibson, D. J. Merenstein, B. Pot, L. Morelli, R. B. Canani, H. J. Flint, S. Salminen, P. C. Calder and M. E. Sanders. (2014). Expert consensus document. The International Scientific Association for Probiotics and Prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nat Rev Gastroenterol Hepatol*, 11(8), 506-514. doi:10.1038/nrgastro.2014.66

Hsieh, C. L., C. C. Kim, B. E. Ryba, E. C. Niemi, J. K. Bando, R. M. Locksley, J. Liu, M. C. Nakamura and W. E. Seaman. (2013). Traumatic brain injury induces macrophage subsets in the brain. *Eur J Immunol*, 43(8), 2010-2022. doi:10.1002/eji.201243084

Hsieh, C. L., E. C. Niemi, S. H. Wang, C. C. Lee, D. Bingham, J. Zhang, M. L. Cozen, I. Charo, E. J. Huang, J. Liu and M. C. Nakamura. (2014). CCR2 deficiency impairs macrophage infiltration and improves cognitive function after traumatic brain injury. *J Neurotrauma*, 31(20), 1677-1688. doi:10.1089/neu.2013.3252

Ianiro, G., H. Tilg and A. Gasbarrini. (2016). Antibiotics as deep modulators of gut microbiota: between good and evil. *Gut*, 65(11), 1906-1915. doi:10.1136/gutjnl-2016-312297

Iniguez, S. D., L. M. Riggs, S. J. Nieto, G. Dayrit, N. N. Zamora, K. L. Shawhan, B. Cruz and B. L. Warren. (2014). Social defeat stress induces a depression-like phenotype in adolescent male c57BL/6 mice. *Stress*, 17(3), 247-255. doi:10.3109/10253890.2014.910650

Israelsson, C., H. Bengtsson, A. Kylberg, K. Kullander, A. Lewen, L. Hillered and T. Ebendal. (2008). Distinct cellular patterns of upregulated chemokine expression supporting a prominent inflammatory role in traumatic brain injury. *J Neurotrauma*, 25(8), 959-974. doi:10.1089/neu.2008.0562

Israelsson, C., A. Kylberg, H. Bengtsson, L. Hillered and T. Ebendal. (2014). Interacting chemokine signals regulate dendritic cells in acute brain injury. *PLoS One*, 9(8), e104754. doi:10.1371/journal.pone.0104754

Izikson, L., R. S. Klein, I. F. Charo, H. L. Weiner and A. D. Luster. (2000). Resistance to experimental autoimmune encephalomyelitis in mice lacking the CC chemokine receptor (CCR)2. *J Exp Med*, 192(7), 1075-1080.

Jakubzick, C. V., G. J. Randolph and P. M. Henson. (2017). Monocyte differentiation and antigen-presenting functions. *Nat Rev Immunol*, 17(6), 349-362. doi:10.1038/nri.2017.28

Jeong, J. J., K. A. Kim, Y. T. Ahn, J. H. Sim, J. Y. Woo, C. S. Huh and D. H. Kim. (2015). Probiotic Mixture KF Attenuates Age-Dependent Memory Deficit and Lipidemia in Fischer 344 Rats. *J Microbiol Biotechnol*, 25(9), 1532-1536. doi:10.4014/jmb.1505.05002

Johnson, J. D., K. A. O'Connor, T. Deak, M. Stark, L. R. Watkins and S. F. Maier. (2002). Prior stressor exposure sensitizes LPS-induced cytokine production. *Brain Behav Immun*, 16(4), 461-476.

Jordao, M. J. C., R. Sankowski, S. M. Brendecke, Sagar, G. Locatelli, Y. H. Tai, T. L. Tay, E. Schramm, S. Armbruster, N. Hagemeyer, O. Gross, D. Mai, O. Cicek, T. Falk, M. Kerschensteiner, D. Grun and M. Prinz. (2019). Single-cell profiling identifies myeloid cell subsets with distinct fates during neuroinflammation. *Science*, 363(6425). doi:10.1126/science.aat7554

Kang, D. W., J. B. Adams, D. M. Coleman, E. L. Pollard, J. Maldonado, S. McDonough-Means, J. G. Caporaso and R. Krajmalnik-Brown. (2019). Long-term benefit of Microbiota Transfer Therapy on autism symptoms and gut microbiota. *Sci Rep*, 9(1), 5821. doi:10.1038/s41598-019-42183-0

Karlmarm, K. R., F. Tacke and I. R. Dunay. (2012). Monocytes in health and disease - Minireview. *Eur J Microbiol Immunol (Bp)*, 2(2), 97-102. doi:10.1556/EuJMI.2.2012.2.1

Kaufer, C., C. Chhatbar, S. Broer, I. Walzl, L. Ghita, I. Gerhauser, U. Kalinke and W. Loscher. (2018). Chemokine receptors CCR2 and CX3CR1 regulate viral encephalitis-induced hippocampal damage but not seizures. *Proc Natl Acad Sci U S A*, 115(38), E8929-E8938. doi:10.1073/pnas.1806754115

Kelly, J. R., Y. Borre, O. B. C. E. Patterson, S. El Aidy, J. Deane, P. J. Kennedy, S. Beers, K. Scott, G. Moloney, A. E. Hoban, L. Scott, P. Fitzgerald, P. Ross, C. Stanton, G. Clarke, J. F. Cryan and T. G. Dinan. (2016). Transferring the blues: Depression-associated gut microbiota induces neurobehavioural changes in the rat. *J Psychiatr Res*, 82, 109-118. doi:10.1016/j.jpsychires.2016.07.019

Kelly, K. A., L. T. Michalovicz, J. V. Miller, V. Castranova, D. B. Miller and J. P. O'Callaghan. (2018). Prior exposure to corticosterone markedly enhances and prolongs the neuroinflammatory response to systemic challenge with LPS. *PLoS One*, 13(1), e0190546. doi:10.1371/journal.pone.0190546

Kerfoot, S. M., C. D'Mello, H. Nguyen, M. N. Ajuebor, P. Kubes, T. Le and M. G. Swain. (2006). TNF-alpha-secreting monocytes are recruited into the brain of cholestatic mice. *Hepatology*, 43(1), 154-162. doi:10.1002/hep.21003

Keshavarzian, A., S. J. Green, P. A. Engen, R. M. Voigt, A. Naqib, C. B. Forsyth, E. Mutlu and K. M. Shannon. (2015). Colonic bacterial composition in Parkinson's disease. *Mov Disord*, 30(10), 1351-1360. doi:10.1002/mds.26307

Khosravi, A., A. Yanez, J. G. Price, A. Chow, M. Merad, H. S. Goodridge and S. K. Mazmanian. (2014). Gut microbiota promote hematopoiesis to control bacterial infection. *Cell Host Microbe*, 15(3), 374-381. doi:10.1016/j.chom.2014.02.006

Kim, B., V. M. Hong, J. Yang, H. Hyun, J. J. Im, J. Hwang, S. Yoon and J. E. Kim. (2016). A Review of Fermented Foods with Beneficial Effects on Brain and Cognitive Function. *Prev Nutr Food Sci*, 21(4), 297-309. doi:10.3746/pnf.2016.21.4.297

Kim, H. S., J. T. Hong, Y. Kim and S. B. Han. (2011). Stimulatory Effect of beta-glucans on Immune Cells. *Immune Netw*, 11(4), 191-195. doi:10.4110/in.2011.11.4.191

Kim, K. A., J. J. Jeong, S. Y. Yoo and D. H. Kim. (2016). Gut microbiota lipopolysaccharide accelerates inflamm-aging in mice. *BMC Microbiol*, 16, 9. doi:10.1186/s12866-016-0625-7

Kim, S., S. H. Kwon, T. I. Kam, N. Panicker, S. S. Karuppagounder, S. Lee, J. H. Lee, W. R. Kim, M. Kook, C. A. Foss, C. Shen, H. Lee, S. Kulkarni, P. J. Pasricha, G. Lee, M. G. Pomper, V. L. Dawson, T. M. Dawson and H. S. Ko. (2019). Transneuronal Propagation of Pathologic alpha-Synuclein from the Gut to the Brain Models Parkinson's Disease. *Neuron*. doi:10.1016/j.neuron.2019.05.035

Klaus, F., J. C. Paterna, E. Marzorati, H. Sigrist, L. Gotze, S. Schwendener, G. Bergamini, E. Jehli, D. Azzinnari, R. Fuertig, A. Fontana, E. Seifritz and C. R. Pryce. (2016). Differential effects of peripheral and brain tumor necrosis factor on inflammation, sickness, emotional behavior and memory in mice. *Brain Behav Immun*, 58, 310-326. doi:10.1016/j.bbi.2016.08.001

Koh, A., F. De Vadder, P. Kovatcheva-Datchary and F. Backhed. (2016). From Dietary Fiber to Host Physiology: Short-Chain Fatty Acids as Key Bacterial Metabolites. *Cell*, 165(6), 1332-1345. doi:10.1016/j.cell.2016.05.041

Korin, B., T. L. Ben-Shaanan, M. Schiller, T. Dubovik, H. Azulay-Debby, N. T. Boshnak, T. Koren and A. Rolls. (2017). High-dimensional, single-cell characterization of the brain's immune compartment. *Nat Neurosci*, advance online publication. doi:10.1038/nn.4610

<http://www.nature.com/neuro/journal/vaop/ncurrent/abs/nn.4610.html#supplementary-information>

Kouchaki, E., O. R. Tamtaji, M. Salami, F. Bahmani, R. Daneshvar Kakhaki, E. Akbari, M. Tajabadi-Ebrahimi, P. Jafari and Z. Asemi. (2017). Clinical and metabolic response to probiotic supplementation in patients with multiple sclerosis: A randomized, double-blind, placebo-controlled trial. *Clin Nutr*, 36(5), 1245-1249. doi:10.1016/j.clnu.2016.08.015

Lach, G., H. Schellekens, T. G. Dinan and J. F. Cryan. (2018). Anxiety, Depression, and the Microbiome: A Role for Gut Peptides. *Neurotherapeutics*, 15(1), 36-59. doi:10.1007/s13311-017-0585-0

Lafuse, W. P., R. Gearinger, S. Fisher, C. Nealer, A. R. Mackos and M. T. Bailey. (2017). Exposure to a Social Stressor Induces Translocation of Commensal Lactobacilli to the Spleen and Priming of the Innate Immune System. *J Immunol*, 198(6), 2383-2393. doi:10.4049/jimmunol.1601269

Langdon, A., N. Crook and G. Dantas. (2016). The effects of antibiotics on the microbiome throughout development and alternative approaches for therapeutic modulation. *Genome Med*, 8(1), 39. doi:10.1186/s13073-016-0294-z

Langgartner, D., C. A. Lowry and S. O. Reber. (2018). Old Friends, immunoregulation, and stress resilience. *Pflugers Arch*. doi:10.1007/s00424-018-2228-7

Langgartner, D., D. Peterlik, S. Foertsch, A. M. Fuchsl, P. Brokmann, P. J. Flor, Z. Shen, J. G. Fox, N. Uschold-Schmidt, C. A. Lowry and S. O. Reber. (2016). Individual differences in stress vulnerability: the role of gut pathobionts in stress-induced colitis. *Brain Behav Immun*. doi:10.1016/j.bbi.2016.12.019

Lasitschka, F., T. Giese, M. Paparella, S. R. Kurzhals, G. Wabnitz, K. Jacob, J. Gras, K. A. Bode, A. K. Heninger, T. Sziskai, Y. Samstag, C. Leszinski, B. Jocher, M. Al-Saeedi, S. C. Meuer and J. Schroder-Braunstein. (2017). Human monocytes downregulate innate response receptors following exposure to the microbial metabolite n-butyrate. *Immun Inflamm Dis*. doi:10.1002/iid3.184

Lehmann, M. L., H. A. Cooper, D. Maric and M. Herkenham. (2016). Social defeat induces depressive-like states and microglial activation without involvement of peripheral macrophages. *J Neuroinflammation*, 13(1), 224. doi:10.1186/s12974-016-0672-x

Lehmann, M. L., T. K. Weigel, H. A. Cooper, A. G. Elkahloun, S. L. Kigar and M. Herkenham. (2018). Decoding microglia responses to psychosocial stress reveals blood-brain barrier breakdown that may drive stress susceptibility. *Sci Rep*, 8(1), 11240. doi:10.1038/s41598-018-28737-8

Li, H., J. Sun, J. Du, F. Wang, R. Fang, C. Yu, J. Xiong, W. Chen, Z. Lu and J. Liu. (2018). Clostridium butyricum exerts a neuroprotective effect in a mouse model of traumatic brain injury via the gut-brain axis. *Neurogastroenterol Motil*, 30(5), e13260. doi:10.1111/nmo.13260

Li, J., L. Hou, C. Wang, X. Jia, X. Qin and C. Wu. (2018). Short Term Intrarectal Administration of Sodium Propionate Induces Antidepressant-Like Effects in Rats Exposed to Chronic Unpredictable Mild Stress. *Front Psychiatry*, 9, 454. doi:10.3389/fpsy.2018.00454

Li, M., B. van Esch, G. T. M. Wagenaar, J. Garssen, G. Folkerts and P. A. J. Henricks. (2018). Pro- and anti-inflammatory effects of short chain fatty acids on immune and endothelial cells. *Eur J Pharmacol*. doi:10.1016/j.ejphar.2018.05.003

Li, N., Q. Wang, Y. Wang, A. Sun, Y. Lin, Y. Jin and X. Li. (2018). Oral Probiotics Ameliorate the Behavioral Deficits Induced by Chronic Mild Stress in Mice via the Gut Microbiota-Inflammation Axis. *Front Behav Neurosci*, 12, 266. doi:10.3389/fnbeh.2018.00266

Liang, S., T. Wang, X. Hu, J. Luo, W. Li, X. Wu, Y. Duan and F. Jin. (2015). Administration of Lactobacillus helveticus NS8 improves behavioral, cognitive, and biochemical aberrations caused by chronic restraint stress. *Neuroscience*, 310, 561-577. doi:10.1016/j.neuroscience.2015.09.033

Lisboa, S. F., A. Niraula, L. B. Resstel, F. S. Guimaraes, J. P. Godbout and J. F. Sheridan. (2018). Repeated social defeat-induced neuroinflammation, anxiety-like behavior and resistance to fear extinction were attenuated by the cannabinoid receptor agonist WIN55,212-2. *Neuropsychopharmacology*. doi:10.1038/s41386-018-0064-2

Liu, B., F. Fang, N. L. Pedersen, A. Tillander, J. F. Ludvigsson, A. Ekbom, P. Svenningsson, H. Chen and K. Wirdefeldt. (2017). Vagotomy and Parkinson disease: A Swedish register-based matched-cohort study. *Neurology*, 88(21), 1996-2002. doi:10.1212/WNL.0000000000003961

Liu, K., G. D. Victora, T. A. Schwickert, P. Guermonprez, M. M. Meredith, K. Yao, F. F. Chu, G. J. Randolph, A. Y. Rudensky and M. Nussenzweig. (2009). In vivo analysis of dendritic cell development and homeostasis. *Science*, 324(5925), 392-397. doi:10.1126/science.1170540

Loane, D. J. and A. Kumar. (2016). Microglia in the TBI brain: The good, the bad, and the dysregulated. *Exp Neurol*, 275 Pt 3, 316-327. doi:10.1016/j.expneurol.2015.08.018

long-Smith, C. M., K. J. O'Riordan, G. clarke, C. Stanton, D. T. G. and J. F. Cryan. (2019). Microbiota-Gut-Brain Axis: New Therapeutic Opportunities. *Annual Review of Pharmacology and Toxicology*, 60.

Lund, H., M. Pieber, R. Parsa, J. Han, D. Grommisch, E. Ewing, L. Kular, M. Needhamsen, A. Espinosa, E. Nilsson, A. K. Overby, O. Butovsky, M. Jagodic, X. M. Zhang and R. A. Harris. (2018). Competitive repopulation of an empty microglial niche yields functionally distinct subsets of microglia-like cells. *Nat Commun*, 9(1), 4845. doi:10.1038/s41467-018-07295-7

Luo, Y., B. Zeng, L. Zeng, X. Du, B. Li, R. Huo, L. Liu, H. Wang, M. Dong, J. Pan, P. Zheng, C. Zhou, H. Wei and P. Xie. (2018). Gut microbiota regulates mouse behaviors through glucocorticoid receptor pathway genes in the hippocampus. *Transl Psychiatry*, 8(1), 187. doi:10.1038/s41398-018-0240-5

Luthold, R. V., G. R. Fernandes, A. C. Franco-de-Moraes, L. G. Folchetti and S. R. Ferreira. (2017). Gut microbiota interactions with the immunomodulatory role of vitamin D in normal individuals. *Metabolism*, 69, 76-86. doi:10.1016/j.metabol.2017.01.007

Ma, Y., T. Liu, J. Fu, S. Fu, C. Hu, B. Sun, X. Fan and J. Zhu. (2019). Lactobacillus acidophilus Exerts Neuroprotective Effects in Mice with Traumatic Brain Injury. *J Nutr*. doi:10.1093/jn/nxz105

Macfarlane, S. and G. T. Macfarlane. (2003). Regulation of short-chain fatty acid production. *Proc Nutr Soc*, 62(1), 67-72. doi:10.1079/PNS2002207

Mackos, A. R., J. D. Galley, T. D. Eubank, R. S. Easterling, N. M. Parry, J. G. Fox, M. Lyte and M. T. Bailey. (2016). Social stress-enhanced severity of Citrobacter rodentium-induced colitis is CCL2-dependent and attenuated by probiotic Lactobacillus reuteri. *Mucosal Immunol*, 9(2), 515-526. doi:10.1038/mi.2015.81

Maehata, H., Y. Kobayashi, E. Mitsuyama, T. Kawase, T. Kuhara, J. Z. Xiao, T. Tsukahara and A. Toyoda. (2019). Heat-killed Lactobacillus helveticus strain MCC1848 confers resilience to anxiety or depression-like symptoms caused by subchronic social defeat stress in mice. *Biosci Biotechnol Biochem*, 1-9. doi:10.1080/09168451.2019.1591263

Makinde, H. M., C. M. Cuda, T. B. Just, H. R. Perlman and S. J. Schwulst. (2017). Nonclassical Monocytes Mediate Secondary Injury, Neurocognitive Outcome, and Neutrophil Infiltration after Traumatic Brain Injury. *J Immunol*, 199(10), 3583-3591. doi:10.4049/jimmunol.1700896

Makinde, H. M., T. B. Just, C. M. Cuda, N. Bertolino, D. Procissi and S. J. Schwulst. (2018). Monocyte depletion attenuates the development of posttraumatic hydrocephalus and preserves white matter integrity after traumatic brain injury. *PLoS One*, 13(11), e0202722. doi:10.1371/journal.pone.0202722

Marin, I. A., J. E. Goertz, T. Ren, S. S. Rich, S. Onengut-Gumuscu, E. Farber, M. Wu, C. C. Overall, J. Kipnis and A. Gaultier. (2017). Microbiota alteration is associated with the development of stress-induced despair behavior. *Sci Rep*, 7, 43859. doi:10.1038/srep43859

Markowiak, P. and K. Slizewska. (2017). Effects of Probiotics, Prebiotics, and Synbiotics on Human Health. *Nutrients*, 9(9). doi:10.3390/nu9091021

Marsland, A. L., C. Walsh, K. Lockwood and N. A. John-Henderson. (2017). The effects of acute psychological stress on circulating and stimulated inflammatory markers: A systematic review and meta-analysis. *Brain Behav Immun*, 64, 208-219. doi:10.1016/j.bbi.2017.01.011

Martin, E., C. Boucher, B. Fontaine and C. Delarasse. (2017). Distinct inflammatory phenotypes of microglia and monocyte-derived macrophages in Alzheimer's disease models: effects of aging and amyloid pathology. *Aging Cell*, 16(1), 27-38. doi:10.1111/acel.12522

Martinez, L., C. Gomez and R. I. Vazquez-Padron. (2015). Age-related changes in monocytes exacerbate neointimal hyperplasia after vascular injury. *Oncotarget*, 6(19), 17054-17064. doi:10.18632/oncotarget.3881

Matt, S. M., J. M. Allen, M. A. Lawson, L. J. Mailing, J. A. Woods and R. W. Johnson. (2018). Butyrate and Dietary Soluble Fiber Improve Neuroinflammation Associated With Aging in Mice. *Front Immunol*, 9, 1832. doi:10.3389/fimmu.2018.01832

McEwen, B. S., J. D. Gray and C. Nasca. (2015). 60 YEARS OF NEUROENDOCRINOLOGY: Redefining neuroendocrinology: stress, sex and cognitive and emotional regulation. *J Endocrinol*, 226(2), T67-83. doi:10.1530/JOE-15-0121

McKim, D. B., J. M. Patterson, E. S. Wohleb, B. L. Jarrett, B. F. Reader, J. P. Godbout and J. F. Sheridan. (2016). Sympathetic Release of Splenic Monocytes Promotes Recurring Anxiety Following Repeated Social Defeat. *Biol Psychiatry*, 79(10), 803-813. doi:10.1016/j.biopsych.2015.07.010

McKim, D. B., W. Yin, Y. Wang, S. W. Cole, J. P. Godbout and J. F. Sheridan. (2018). Social Stress Mobilizes Hematopoietic Stem Cells to Establish Persistent Splenic Myelopoiesis. *Cell Rep*, 25(9), 2552-2562 e2553. doi:10.1016/j.celrep.2018.10.102

Mestre, L., F. J. Carrillo-Salinas, M. Mecha, A. Feliu, C. Espejo, J. C. Alvarez-Cermeno, L. M. Villar and C. Guaza. (2019). Manipulation of Gut Microbiota Influences Immune Responses, Axon Preservation, and Motor Disability in a Model of Progressive Multiple Sclerosis. *Front Immunol*, 10, 1374. doi:10.3389/fimmu.2019.01374

Michaud, J. P., M. A. Bellavance, P. Prefontaine and S. Rivest. (2013). Real-time in vivo imaging reveals the ability of monocytes to clear vascular amyloid beta. *Cell Rep*, 5(3), 646-653. doi:10.1016/j.celrep.2013.10.010

Mika, A., H. E. Day, A. Martinez, N. L. Rumian, B. N. Greenwood, M. Chichlowski, B. M. Berg and M. Fleshner. (2017). Early life diets with prebiotics and bioactive milk fractions attenuate the impact of stress on learned helplessness behaviours and alter gene expression within neural circuits important for stress resistance. *Eur J Neurosci*, 45(3), 342-357. doi:10.1111/ejn.13444

Mildner, A., M. Mack, H. Schmidt, W. Bruck, M. Djukic, M. D. Zabel, A. Hille, J. Priller and M. Prinz. (2009). CCR2+Ly-6Chi monocytes are crucial for the effector phase of autoimmunity in the central nervous system. *Brain*, 132(Pt 9), 2487-2500. doi:10.1093/brain/awp144

Mildner, A., B. Schlevogt, K. Kierdorf, C. Bottcher, D. Erny, M. P. Kummer, M. Quinn, W. Bruck, I. Bechmann, M. T. Heneka, J. Priller and M. Prinz. (2011). Distinct and non-redundant roles of microglia and myeloid subsets in mouse models of Alzheimer's disease. *J Neurosci*, 31(31), 11159-11171. doi:10.1523/JNEUROSCI.6209-10.2011

Miller, A. H., R. L. Spencer, B. D. Pearce, T. L. Pisell, Y. Azrieli, P. Tanapat, H. Moday, R. Rhee and B. S. McEwen. (1998). Glucocorticoid receptors are differentially expressed in the cells and tissues of the immune system. *Cell Immunol*, 186(1), 45-54. doi:10.1006/cimm.1998.1293

Miller, G. E., S. Cohen and A. K. Ritchey. (2002). Chronic psychological stress and the regulation of pro-inflammatory cytokines: a glucocorticoid-resistance model. *Health Psychol*, 21(6), 531-541.

Minter, M. R., C. Zhang, V. Leone, D. L. Ringus, X. Zhang, P. Oyler-Castrillo, M. W. Musch, F. Liao, J. F. Ward, D. M. Holtzman, E. B. Chang, R. E. Tanzi and S. S. Sisodia. (2016). Antibiotic-induced perturbations in gut microbial diversity influences neuro-inflammation and amyloidosis in a murine model of Alzheimer's disease. *Sci Rep*, 6, 30028. doi:10.1038/srep30028

Mohle, L., D. Mattei, M. M. Heimesaat, S. Bereswill, A. Fischer, M. Alutis, T. French, D. Hambardzumyan, P. Matzinger, I. R. Dunay and S. A. Wolf. (2016). Ly6C(hi) Monocytes Provide a Link between Antibiotic-Induced Changes in Gut Microbiota and Adult Hippocampal Neurogenesis. *Cell Rep*, 15(9), 1945-1956. doi:10.1016/j.celrep.2016.04.074

Möhle, L., D. Mattei, Markus M. Heimesaat, S. Bereswill, A. Fischer, M. Alutis, T. French, D. Hambardzumyan, P. Matzinger, Ildiko R. Dunay and Susanne A. Wolf. (2016). Ly6Chi Monocytes Provide a Link between Antibiotic-Induced Changes in Gut Microbiota and Adult Hippocampal Neurogenesis. *Cell Reports*, 15(9), 1945-1956. doi:<http://dx.doi.org/10.1016/j.celrep.2016.04.074>

Morganti, J. M., T. D. Jopson, S. Liu, L. K. Riparip, C. K. Guandique, N. Gupta, A. R. Ferguson and S. Rosi. (2015). CCR2 antagonism alters brain macrophage polarization and ameliorates cognitive dysfunction induced by traumatic brain injury. *J Neurosci*, 35(2), 748-760. doi:10.1523/JNEUROSCI.2405-14.2015

Moskowitz, M. A., E. H. Lo and C. Iadecola. (2010). The science of stroke: mechanisms in search of treatments. *Neuron*, 67(2), 181-198. doi:10.1016/j.neuron.2010.07.002

Most, J., J. Penders, M. Lucchesi, G. H. Goossens and E. E. Blaak. (2017). Gut microbiota composition in relation to the metabolic response to 12-week combined polyphenol supplementation in overweight men and women. *Eur J Clin Nutr*, 71(9), 1040-1045. doi:10.1038/ejcn.2017.89

Mrdjen, D., A. Pavlovic, F. J. Hartmann, B. Schreiner, S. G. Utz, B. P. Leung, I. Lelios, F. L. Heppner, J. Kipnis, D. Merkler, M. Greter and B. Becher. (2018). High-Dimensional Single-Cell Mapping of Central Nervous System Immune Cells Reveals Distinct Myeloid Subsets in Health, Aging, and Disease. *Immunity*, 48(3), 599. doi:10.1016/j.immuni.2018.02.014

Naert, G. and S. Rivest. (2011). CC chemokine receptor 2 deficiency aggravates cognitive impairments and amyloid pathology in a transgenic mouse model of Alzheimer's disease. *J Neurosci*, 31(16), 6208-6220. doi:10.1523/JNEUROSCI.0299-11.2011

Naert, G. and S. Rivest. (2012). Hematopoietic CC-chemokine receptor 2 (CCR2) competent cells are protective for the cognitive impairments and amyloid pathology in a transgenic mouse model of Alzheimer's disease. *Mol Med*, 18, 297-313. doi:10.2119/molmed.2011.00306

Nie, X., S. Kitaoka, K. Tanaka, E. Segi-Nishida, Y. Imoto, A. Ogawa, F. Nakano, A. Tomohiro, K. Nakayama, M. Taniguchi, Y. Mimori-Kiyosue, A. Kakizuka, S. Narumiya and T. Furuyashiki. (2018). The Innate Immune Receptors TLR2/4 Mediate Repeated Social Defeat Stress-Induced Social Avoidance through Prefrontal Microglial Activation. *Neuron*, 99(3), 464-479 e467. doi:10.1016/j.neuron.2018.06.035

Niraula, A., Y. Wang, J. P. Godbout and J. F. Sheridan. (2018). Corticosterone Production during Repeated Social Defeat Causes Monocyte Mobilization from the Bone Marrow, Glucocorticoid Resistance and Neurovascular Adhesion Molecule Expression. *J Neurosci*. doi:10.1523/JNEUROSCI.2568-17.2018

Niraula, A., K. G. Witcher, J. F. Sheridan and J. P. Godbout. (2018). Interleukin-6 Induced by Social Stress Promotes a Unique Transcriptional Signature in the Monocytes That Facilitate Anxiety. *Biol Psychiatry*. doi:10.1016/j.biopsych.2018.09.030

Ohsawa, K., F. Nakamura, N. Uchida, S. Mizuno and H. Yokogoshi. (2018). Lactobacillus helveticus-fermented milk containing lactononadecapeptide (NIPPLTQTPVVPPFLQPE) improves cognitive function in healthy middle-aged adults: a randomised, double-blind, placebo-controlled trial. *Int J Food Sci Nutr*, 69(3), 369-376. doi:10.1080/09637486.2017.1365824

Oishi, Y. and I. Manabe. (2016). Macrophages in age-related chronic inflammatory diseases. *NPJ Aging Mech Dis*, 2, 16018. doi:10.1038/npjamd.2016.18

Okutsu, M., K. Suzuki, T. Ishijima, J. Peake and M. Higuchi. (2008). The effects of acute exercise-induced cortisol on CCR2 expression on human monocytes. *Brain Behav Immun*, 22(7), 1066-1071. doi:10.1016/j.bbi.2008.03.006

Olnes, M. J., Y. Kotliarov, A. Biancotto, F. Cheung, J. Chen, R. Shi, H. Zhou, E. Wang, J. S. Tsang, R. Nussenblatt and C. H. I. Consortium. (2016). Effects of Systemically Administered Hydrocortisone on the Human Immunome. *Sci Rep*, 6, 23002. doi:10.1038/srep23002

Östlund-Lagerström, L., A. Kihlgren, D. Repsilber, B. Björkstén, R. J. Brummer and I. Schoultz. (2016). Probiotic administration among free-living older adults: a double blinded, randomized, placebo-controlled clinical trial. *Nutr J*, 15(1), 80. doi:10.1186/s12937-016-0198-1

Palframan, R. T., S. Jung, G. Cheng, W. Weninger, Y. Luo, M. Dorf, D. R. Littman, B. J. Rollins, H. Zweerink, A. Rot and U. H. von Andrian. (2001). Inflammatory chemokine transport and presentation in HEV: a remote control mechanism for monocyte recruitment to lymph nodes in inflamed tissues. *J Exp Med*, 194(9), 1361-1373.

Palin, K., R. M. Bluthé, R. H. McCusker, F. Moos, R. Dantzer and K. W. Kelley. (2007). TNF α -induced sickness behavior in mice with functional 55 kD TNF receptors is blocked by central IGF-I. *J Neuroimmunol*, 187(1-2), 55-60. doi:10.1016/j.jneuroim.2007.04.011

Paradis, A., S. Bernier and N. Dumais. (2016). TLR4 induces CCR7-dependent monocytes transmigration through the blood-brain barrier. *J Neuroimmunol*, 295-296, 12-17. doi:10.1016/j.jneuroim.2016.03.019

Passlick, B., D. Flieger and H. W. Ziegler-Heitbrock. (1989). Identification and characterization of a novel monocyte subpopulation in human peripheral blood. *Blood*, 74(7), 2527-2534.

Patel, A. A., Y. Zhang, J. N. Fullerton, L. Boelen, A. Rongvaux, A. A. Maini, V. Bigley, R. A. Flavell, D. W. Gilroy, B. Asquith, D. Macallan and S. Yona. (2017). The fate and lifespan of human monocyte subsets in steady state and systemic inflammation. *J Exp Med*, 214(7), 1913-1923. doi:10.1084/jem.20170355

Penton-Rol, G., M. Cota, N. Polentarutti, W. Luini, S. Bernasconi, A. Borsatti, A. Sica, G. J. LaRosa, S. Sozzani, G. Poli and A. Mantovani. (1999). Up-regulation of CCR2 chemokine receptor expression and increased susceptibility to the multitropic HIV strain 89.6 in monocytes exposed to glucocorticoid hormones. *J Immunol*, 163(6), 3524-3529.

Peterlik, D., C. Stangl, A. Bauer, A. Bludau, J. Keller, D. Grabski, T. Killian, D. Schmidt, F. Zajicek, G. Jaeschke, L. Lindemann, S. O. Reber, P. J. Flor and N. Uschold-Schmidt. (2017). Blocking metabotropic glutamate receptor subtype 5 relieves maladaptive chronic stress consequences. *Brain Behav Immun*, 59, 79-92. doi:10.1016/j.bbi.2016.08.007

Peters, S. G., E. W. Pomare and C. A. Fisher. (1992). Portal and peripheral blood short chain fatty acid concentrations after caecal lactulose instillation at surgery. *Gut*, 33(9), 1249-1252.

Plovier, H. and P. D. Cani. (2017). Enteroendocrine Cells: Metabolic Relays between Microbes and Their Host. *Endocr Dev*, 32, 139-164. doi:10.1159/000475736

Powell, N. D., E. K. Sloan, M. T. Bailey, J. M. Arevalo, G. E. Miller, E. Chen, M. S. Kobor, B. F. Reader, J. F. Sheridan and S. W. Cole. (2013). Social stress up-regulates inflammatory gene expression in the leukocyte transcriptome via beta-adrenergic induction of myelopoiesis. *Proc Natl Acad Sci U S A*, 110(41), 16574-16579. doi:10.1073/pnas.1310655110

Prinz, M. and J. Priller. (2010). Tickets to the brain: role of CCR2 and CX3CR1 in myeloid cell entry in the CNS. *J Neuroimmunol*, 224(1-2), 80-84. doi:10.1016/j.jneuroim.2010.05.015

Prinz, M., J. Priller, S. S. Sisodia and R. M. Ransohoff. (2011). Heterogeneity of CNS myeloid cells and their roles in neurodegeneration. *Nat Neurosci*, 14(10), 1227-1235. doi:10.1038/nn.2923

Provinsi, G., S. D. Schmidt, M. Boehme, T. F. S. Bastiaanssen, B. Rani, A. Costa, K. Busca, F. Fouhy, C. Strain, C. Stanton, P. Blandina, I. Izquierdo, J. F. Cryan and M. B. Passani. (2019). Preventing adolescent stress-induced cognitive and microbiome changes by diet. *Proc Natl Acad Sci U S A*, 116(19), 9644-9651. doi:10.1073/pnas.1820832116

Puchta, A., A. Naidoo, C. P. Verschoor, D. Loukov, N. Thevaranjan, T. S. Mandur, P. S. Nguyen, M. Jordana, M. Loeb, Z. Xing, L. Kobzik, M. J. Larche and D. M. Bowdish. (2016). TNF Drives Monocyte Dysfunction with Age and Results in Impaired Anti-pneumococcal Immunity. *PLoS Pathog*, 12(1), e1005368. doi:10.1371/journal.ppat.1005368

Pusceddu, M. M., S. El Aidy, F. Crispie, O. O'Sullivan, P. Cotter, C. Stanton, P. Kelly, J. F. Cryan and T. G. Dinan. (2015). N-3 Polyunsaturated Fatty Acids (PUFAs) Reverse the Impact of Early-Life Stress on the Gut Microbiota. *PLoS One*, 10(10), e0139721. doi:10.1371/journal.pone.0139721

Quintin, J., S. Saeed, J. H. A. Martens, E. J. Giamarellos-Bourboulis, D. C. Ifrim, C. Logie, L. Jacobs, T. Jansen, B. J. Kullberg, C. Wijmenga, L. A. B. Joosten, R. J. Xavier, J. W. M. van der Meer, H. G. Stunnenberg and M. G. Netea. (2012). *Candida albicans* infection affords protection against reinfection via functional reprogramming of monocytes. *Cell Host Microbe*, 12(2), 223-232. doi:10.1016/j.chom.2012.06.006

Ramirez, K., J. Fornaguera-Trias and J. F. Sheridan. (2017). Stress-Induced Microglia Activation and Monocyte Trafficking to the Brain Underlie the Development of Anxiety and Depression. *Curr Top Behav Neurosci*, 31, 155-172. doi:10.1007/7854_2016_25

Ransohoff, R. M. (2011). Microglia and monocytes: 'tis plain the twain meet in the brain. *Nature Neuroscience*, 14, 1098. doi:10.1038/nn.2917

Ransohoff, R. M. and A. E. Cardona. (2010). The myeloid cells of the central nervous system parenchyma. *Nature*, 468, 253. doi:10.1038/nature09615

Rea, K., T. G. Dinan and J. F. Cryan. (2016). The microbiome: A key regulator of stress and neuroinflammation. *Neurobiol Stress*, 4, 23-33. doi:10.1016/j.ynstr.2016.03.001

Rhee, S. H., C. Pothoulakis and E. A. Mayer. (2009). Principles and clinical implications of the brain-gut-enteric microbiota axis. *Nat Rev Gastroenterol Hepatol*, 6(5), 306-314. doi:10.1038/nrgastro.2009.35

Rhodes, J. K., J. Sharkey and P. J. Andrews. (2009). The temporal expression, cellular localization, and inhibition of the chemokines MIP-2 and MCP-1 after traumatic brain injury in the rat. *J Neurotrauma*, 26(4), 507-525. doi:10.1089/neu.2008.0686

Ridaura, V. K., J. J. Faith, F. E. Rey, J. Cheng, A. E. Duncan, A. L. Kau, N. W. Griffin, V. Lombard, B. Henrissat, J. R. Bain, M. J. Muehlbauer, O. Ilkayeva, C. F. Semenkovich, K. Funai, D. K. Hayashi, B. J. Lyle, M. C. Martini, L. K. Ursell, J. C. Clemente, W. Van Treuren, W. A. Walters, R. Knight, C. B. Newgard, A. C. Heath and J. I. Gordon. (2013). Gut microbiota from twins discordant for obesity modulate metabolism in mice. *Science*, 341(6150), 1241214. doi:10.1126/science.1241214

Rieder, R., P. J. Wisniewski, B. L. Alderman and S. C. Campbell. (2017). Microbes and mental health: A review. *Brain Behav Immun*, 66, 9-17. doi:10.1016/j.bbi.2017.01.016

Robertson, R. C., C. Seira Oriach, K. Murphy, G. M. Moloney, J. F. Cryan, T. G. Dinan, R. Paul Ross and C. Stanton. (2017). Omega-3 polyunsaturated fatty acids critically regulate behaviour and gut microbiota development in adolescence and adulthood. *Brain Behav Immun*, 59, 21-37. doi:10.1016/j.bbi.2016.07.145

Rodriguez-Arias, M., S. Montagud-Romero, A. Rubio-Araiz, M. A. Aguilar, E. Martin-Garcia, R. Cabrera, R. Maldonado, F. Porcu, M. I. Colado and J. Minarro. (2015). Effects of repeated social defeat on adolescent mice on cocaine-induced CPP and self-administration in adulthood: integrity of the blood-brain barrier. *Addict Biol*. doi:10.1111/adb.12301

Rohleder, N. (2014). Stimulation of systemic low-grade inflammation by psychosocial stress. *Psychosom Med*, 76(3), 181-189. doi:10.1097/PSY.0000000000000049

Rooks, M. G. and W. S. Garrett. (2016). Gut microbiota, metabolites and host immunity. *Nat Rev Immunol*, 16(6), 341-352. doi:10.1038/nri.2016.42

Rosadini, C. V. and J. C. Kagan. (2017). Early innate immune responses to bacterial LPS. *Curr Opin Immunol*, 44, 14-19. doi:10.1016/j.coi.2016.10.005

Rossol, M., H. Heine, U. Meusch, D. Quandt, C. Klein, M. J. Sweet and S. Hauschildt. (2011). LPS-induced cytokine production in human monocytes and macrophages. *Crit Rev Immunol*, 31(5), 379-446.

Saeed, S., J. Quintin, H. H. Kerstens, N. A. Rao, A. Aghajani-Refah, F. Matarese, S. C. Cheng, J. Ratter, K. Berentsen, M. A. van der Ent, N. Sharifi, E. M. Janssen-Megens, M. Ter Huurne, A. Mandoli, T. van Schaik, A. Ng, F. Burden, K. Downes, M. Frontini, V. Kumar, E. J. Giamarellos-Bourboulis, W. H. Ouwehand, J. W. van der Meer, L. A. Joosten, C. Wijmenga, J. H. Martens, R. J. Xavier, C. Logie, M. G. Netea and H. G. Stunnenberg. (2014). Epigenetic programming of monocyte-to-macrophage differentiation and trained innate immunity. *Science*, 345(6204), 1251086. doi:10.1126/science.1251086

Saemann, M. D., G. A. Bohmig, C. H. Osterreicher, H. Burtscher, O. Parolini, C. Diakos, J. Stockl, W. H. Horl and G. J. Zlabinger. (2000). Anti-inflammatory effects of sodium butyrate on human monocytes: potent inhibition of IL-12 and up-regulation of IL-10 production. *FASEB J*, 14(15), 2380-2382. doi:10.1096/fj.00-0359fje

Salam, A. P., A. Borsini and P. A. Zunszain. (2018). Trained innate immunity: a salient factor in the pathogenesis of neuroimmune psychiatric disorders. *Mol Psychiatry*, 23(2), 170-176. doi:10.1038/mp.2017.186

Sampson, T. R., J. W. Debelius, T. Thron, S. Janssen, G. G. Shastri, Z. E. Ilhan, C. Challis, C. E. Schretter, S. Rocha, V. Gradinaru, M. F. Chesselet, A. Keshavarzian, K. M. Shannon, R. Krajmalnik-Brown, P. Wittung-Stafshede, R. Knight and S. K. Mazmanian. (2016). Gut Microbiota Regulate Motor Deficits and Neuroinflammation in a Model of Parkinson's Disease. *Cell*, 167(6), 1469-1480 e1412. doi:10.1016/j.cell.2016.11.018

Sandhu, K. V., E. Sherwin, H. Schellekens, C. Stanton, T. G. Dinan and J. F. Cryan. (2017). Feeding the microbiota-gut-brain axis: diet, microbiome, and neuropsychiatry. *Transl Res*, 179, 223-244. doi:10.1016/j.trsl.2016.10.002

Santiago-Tirado, F. H. and T. L. Doering. (2017). False friends: Phagocytes as Trojan horses in microbial brain infections. *PLoS Pathog*, 13(12), e1006680. doi:10.1371/journal.ppat.1006680

Sapolsky, R. M., L. M. Romero and A. U. Munck. (2000). How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocr Rev*, 21(1), 55-89. doi:10.1210/edrv.21.1.0389

Saresella, M., I. Marventano, E. Calabrese, F. Piancone, V. Rainone, A. Gatti, M. Alberoni, R. Nemni and M. Clerici. (2014). A complex proinflammatory role for peripheral monocytes in Alzheimer's disease. *J Alzheimers Dis*, 38(2), 403-413. doi:10.3233/JAD-131160

Sarkar, A., S. M. Lehto, S. Harty, T. G. Dinan, J. F. Cryan and P. W. J. Burnet. (2016). Psychobiotics and the Manipulation of Bacteria-Gut-Brain Signals. *Trends Neurosci*, 39(11), 763-781. doi:10.1016/j.tins.2016.09.002

Savignac, H. M., Y. Couch, M. Stratford, D. M. Bannerman, G. Tzortzis, D. C. Anthony and P. W. J. Burnet. (2016). Prebiotic administration normalizes lipopolysaccharide (LPS)-induced anxiety and cortical 5-HT_{2A} receptor and IL1-beta levels in male mice. *Brain Behav Immun*, 52, 120-131. doi:10.1016/j.bbi.2015.10.007

Sawicki, C. M., D. B. McKim, E. S. Wohleb, B. L. Jarrett, B. F. Reader, D. M. Norden, J. P. Godbout and J. F. Sheridan. (2015). Social defeat promotes a reactive endothelium in a brain region-dependent manner with increased expression of key adhesion molecules, selectins and chemokines associated with the recruitment of myeloid cells to the brain. *Neuroscience*, 302, 151-164. doi:10.1016/j.neuroscience.2014.10.004

Scheperjans, F., V. Aho, P. A. Pereira, K. Koskinen, L. Paulin, E. Pekkonen, E. Haapaniemi, S. Kaakkola, J. Eerola-Rautio, M. Pohja, E. Kinnunen, K. Murros and P. Auvinen. (2015). Gut microbiota are related to Parkinson's disease and clinical phenotype. *Mov Disord*, 30(3), 350-358. doi:10.1002/mds.26069

Schmidt, K., P. J. Cowen, C. J. Harmer, G. Tzortzis, S. Errington and P. W. Burnet. (2015). Prebiotic intake reduces the waking cortisol response and alters emotional bias in healthy volunteers. *Psychopharmacology (Berl)*, 232(10), 1793-1801. doi:10.1007/s00213-014-3810-0

Schulthess, J., S. Pandey, M. Capitani, K. C. Rue-Albrecht, I. Arnold, F. Franchini, A. Chomka, N. E. Iltott, D. G. W. Johnston, E. Pires, J. McCullagh, S. N. Sansom, C. V. Arancibia-Carcamo, H. H. Uhlig and F. Powrie. (2019). The Short Chain Fatty Acid Butyrate Imprints an Antimicrobial Program in Macrophages. *Immunity*. doi:10.1016/j.immuni.2018.12.018

Scott, K. A., M. Ida, V. L. Peterson, J. A. Prenderville, G. M. Moloney, T. Izumo, K. Murphy, A. Murphy, R. P. Ross, C. Stanton, T. G. Dinan and J. F. Cryan. (2017). Revisiting Metchnikoff: Age-related alterations in microbiota-gut-brain axis in the mouse. *Brain Behav Immun*, 65, 20-32. doi:10.1016/j.bbi.2017.02.004

Semple, B. D., N. Bye, M. Rancan, J. M. Ziebell and M. C. Morganti-Kossmann. (2010). Role of CCL2 (MCP-1) in traumatic brain injury (TBI): evidence from severe TBI patients and CCL2-/- mice. *J Cereb Blood Flow Metab*, 30(4), 769-782. doi:10.1038/jcbfm.2009.262

Shaw, A. C., D. R. Goldstein and R. R. Montgomery. (2013). Age-dependent dysregulation of innate immunity. *Nat Rev Immunol*, 13(12), 875-887. doi:10.1038/nri3547

Shen, C., M. T. Chen, X. H. Zhang, X. L. Yin, H. M. Ning, R. Su, H. S. Lin, L. Song, F. Wang, Y. N. Ma, H. L. Zhao, J. Yu and J. W. Zhang. (2016). The PU.1-Modulated MicroRNA-22 Is a Regulator of Monocyte/Macrophage Differentiation and Acute Myeloid Leukemia. *PLoS Genet*, 12(9), e1006259. doi:10.1371/journal.pgen.1006259

Sima, P., L. Vannucci and V. Vetvicka. (2018). beta-glucans and cholesterol (Review). *Int J Mol Med*, 41(4), 1799-1808. doi:10.3892/ijmm.2018.3411

Simard, A. R., D. Soulet, G. Gowing, J. P. Julien and S. Rivest. (2006). Bone marrow-derived microglia play a critical role in restricting senile plaque formation in Alzheimer's disease. *Neuron*, 49(4), 489-502. doi:10.1016/j.neuron.2006.01.022

Singh, V., S. Roth, G. Llovera, R. Sadler, D. Garzetti, B. Stecher, M. Dichgans and A. Liesz. (2016). Microbiota Dysbiosis Controls the Neuroinflammatory Response after Stroke. *J Neurosci*, 36(28), 7428-7440. doi:10.1523/JNEUROSCI.1114-16.2016

Singh, V., R. Sadler, S. Heindl, G. Llovera, S. Roth, C. Benakis and A. Liesz. (2018). The gut microbiome primes a cerebroprotective immune response after stroke. *J Cereb Blood Flow Metab*, 38(8), 1293-1298. doi:10.1177/0271678X18780130

Smits, L. P., K. E. Bouter, W. M. de Vos, T. J. Borody and M. Nieuwdorp. (2013). Therapeutic potential of fecal microbiota transplantation. *Gastroenterology*, 145(5), 946-953. doi:10.1053/j.gastro.2013.08.058

Sorrells, S. F., J. R. Caso, C. D. Munhoz, C. K. Hu, K. V. Tran, Z. D. Miguel, B. Y. Chien and R. M. Sapolsky. (2013). Glucocorticoid signaling in myeloid cells worsens acute CNS injury and inflammation. *J Neurosci*, 33(18), 7877-7889. doi:10.1523/JNEUROSCI.4705-12.2013

Sorrells, S. F. and R. M. Sapolsky. (2010). Glucocorticoids can arm macrophages for innate immune battle. *Brain Behav Immun*, 24(1), 17-18. doi:10.1016/j.bbi.2009.10.004

Spychala, M. S., V. R. Venna, M. Jandzinski, S. J. Doran, D. J. Durgan, B. P. Ganesh, N. J. Ajami, N. Putluri, J. Graf, R. M. Bryan and L. D. McCullough. (2018). Age-related changes in the gut microbiota influence systemic inflammation and stroke outcome. *Ann Neurol*, 84(1), 23-36. doi:10.1002/ana.25250

Stanley, D., L. J. Mason, K. E. Mackin, Y. N. Srikhanta, D. Lyras, M. D. Prakash, K. Nurgali, A. Venegas, M. D. Hill, R. J. Moore and C. H. Wong. (2016). Translocation and dissemination of commensal bacteria in post-stroke infection. *Nat Med*, 22(11), 1277-1284. doi:10.1038/nm.4194

Stanley, D., R. J. Moore and C. H. Y. Wong. (2018). An insight into intestinal mucosal microbiota disruption after stroke. *Sci Rep*, 8(1), 568. doi:10.1038/s41598-017-18904-8

Steer, J. H., D. T. Ma, L. Dusci, G. Garas, K. E. Pedersen and D. A. Joyce. (1998). Altered leucocyte trafficking and suppressed tumour necrosis factor alpha release from peripheral blood monocytes after intra-articular glucocorticoid treatment. *Ann Rheum Dis*, 57(12), 732-737.

Stephen, A. M., M. M. Champ, S. J. Cloran, M. Fleith, L. van Lieshout, H. Mejbörn and V. J. Burley. (2017). Dietary fibre in Europe: current state of knowledge on definitions, sources, recommendations, intakes and relationships to health. *Nutr Res Rev*, 30(2), 149-190. doi:10.1017/S095442241700004X

Stilling, R. M., M. van de Wouw, G. Clarke, C. Stanton, T. G. Dinan and J. F. Cryan. (2016). The neuropharmacology of butyrate: The bread and butter of the microbiota-gut-brain axis? *Neurochem Int*, 99, 110-132. doi:10.1016/j.neuint.2016.06.011

Sudo, N., Y. Chida, Y. Aiba, J. Sonoda, N. Oyama, X. N. Yu, C. Kubo and Y. Koga. (2004). Postnatal microbial colonization programs the hypothalamic-pituitary-adrenal system for stress response in mice. *J Physiol*, 558(Pt 1), 263-275. doi:10.1113/jphysiol.2004.063388

Sun, B. L., W. W. Li, J. Wang, Y. L. Xu, H. L. Sun, D. Y. Tian, Y. J. Wang and X. Q. Yao. (2019). Gut Microbiota Alteration and Its Time Course in a Tauopathy Mouse Model. *J Alzheimers Dis*. doi:10.3233/JAD-181220

Sun, M., R. D. Brady, P. M. Casillas-Espinosa, D. K. Wright, B. D. Semple, H. A. Kim, R. Mychasiuk, C. G. Sobey, T. J. O'Brien, A. Vinh, S. J. McDonald and S. R. Shultz. (2019). Aged rats have an altered immune response and worse outcomes after traumatic brain injury. *Brain Behav Immun*. doi:10.1016/j.bbi.2019.04.038

Sun, Y., W. Geng, Y. Pan, J. Wang, P. Xiao and Y. Wang. (2019). Supplementation with *Lactobacillus kefirifaciens* ZW3 from Tibetan Kefir improves depression-like behavior in stressed mice by modulating the gut microbiota. *Food Funct*, 10(2), 925-937. doi:10.1039/c8fo02096e

Sung, M. M., T. T. Kim, E. Denou, C. M. Soltys, S. M. Hamza, N. J. Byrne, G. Masson, H. Park, D. S. Wishart, K. L. Madsen, J. D. Schertzer and J. R. Dyck. (2017). Improved Glucose Homeostasis in Obese Mice Treated With Resveratrol Is Associated With Alterations in the Gut Microbiome. *Diabetes*, 66(2), 418-425. doi:10.2337/db16-0680

Svensson, E., E. Horvath-Puho, R. W. Thomsen, J. C. Djurhuus, L. Pedersen, P. Borghammer and H. T. Sørensen. (2015). Vagotomy and subsequent risk of Parkinson's disease. *Ann Neurol*, 78(4), 522-529. doi:10.1002/ana.24448

Szmydynger-Chodobska, J., N. Strazielle, J. R. Gandy, T. H. Keefe, B. J. Zink, J. F. Gherzi-Egea and A. Chodobski. (2012). Posttraumatic invasion of monocytes across the blood-cerebrospinal fluid barrier. *J Cereb Blood Flow Metab*, 32(1), 93-104. doi:10.1038/jcbfm.2011.111

Szyszkowicz, J. K., A. Wong, H. Anisman, Z. Merali and M. C. Audet. (2017). Implications of the gut microbiota in vulnerability to the social avoidance effects of chronic social defeat in male mice. *Brain Behav Immun*, 66, 45-55. doi:10.1016/j.bbi.2017.06.009

Takada, M., K. Nishida, A. Kataoka-Kato, Y. Gondo, H. Ishikawa, K. Suda, M. Kawai, R. Hoshi, O. Watanabe, T. Igarashi, Y. Kuwano, K. Miyazaki and K. Rokutan. (2016). Probiotic *Lactobacillus casei* strain Shirota relieves stress-associated symptoms by modulating the gut-brain interaction in human and animal models. *Neurogastroenterol Motil*, 28(7), 1027-1036. doi:10.1111/nmo.12804

Takeshita, Y. and R. M. Ransohoff. (2012). Inflammatory cell trafficking across the blood-brain barrier: chemokine regulation and in vitro models. *Immunol Rev*, 248(1), 228-239. doi:10.1111/j.1600-065X.2012.01127.x

Tan, M., J. C. Zhu, J. Du, L. M. Zhang and H. H. Yin. (2011). Effects of probiotics on serum levels of Th1/Th2 cytokine and clinical outcomes in severe traumatic brain-injured patients: a prospective randomized pilot study. *Crit Care*, 15(6), R290. doi:10.1186/cc10579

Tankou, S. K., K. Regev, B. C. Healy, E. Tjon, L. Laghi, L. M. Cox, P. Kivisakk, I. V. Pierre, L. Hrishikesh, R. Gandhi, S. Cook, B. Glanz, J. Stankiewicz and H. L. Weiner. (2018). A probiotic modulates the microbiome and immunity in multiple sclerosis. *Ann Neurol*, 83(6), 1147-1161. doi:10.1002/ana.25244

Tarr, A. J., J. D. Galley, S. E. Fisher, M. Chichlowski, B. M. Berg and M. T. Bailey. (2015). The prebiotics 3'Sialyllactose and 6'Sialyllactose diminish stressor-induced anxiety-like behavior and colonic microbiota alterations: Evidence for effects on the gut-brain axis. *Brain Behav Immun*, 50, 166-177. doi:10.1016/j.bbi.2015.06.025

Thevaranjan, N., A. Puchta, C. Schulz, A. Naidoo, J. C. Szamosi, C. P. Verschoor, D. Loukov, L. P. Schenck, J. Jury, K. P. Foley, J. D. Schertzer, M. J. Larche, D. J. Davidson, E. F. Verdu, M. G. Surette and D. M. E. Bowdish. (2017). Age-Associated Microbial Dysbiosis Promotes Intestinal Permeability, Systemic Inflammation, and Macrophage Dysfunction. *Cell Host Microbe*, 21(4), 455-466 e454. doi:10.1016/j.chom.2017.03.002

Thome, A. D., A. Faridar, D. R. Beers, J. R. Thonhoff, W. Zhao, S. Wen, B. Pascual, J. C. Masdeu and S. H. Appel. (2018). Functional alterations of myeloid cells during the course of Alzheimer's disease. *Mol Neurodegener*, 13(1), 61. doi:10.1186/s13024-018-0293-1

Treangen, T. J., J. Wagner, M. P. Burns and S. Villapol. (2018). Traumatic Brain Injury in Mice Induces Acute Bacterial Dysbiosis Within the Fecal Microbiome. *Front Immunol*, 9, 2757. doi:10.3389/fimmu.2018.02757

Tsianakas, A., G. Varga, K. Barczyk, G. Bode, N. Nippe, N. Kran, J. Roth, T. A. Luger, J. Ehrchen and C. Sunderkoetter. (2012). Induction of an anti-inflammatory human monocyte subtype is a unique property of glucocorticoids, but can be modified by IL-6 and IL-10. *Immunobiology*, 217(3), 329-335. doi:10.1016/j.imbio.2011.10.002

Tsou, C. L., W. Peters, Y. Si, S. Slaymaker, A. M. Aslanian, S. P. Weisberg, M. Mack and I. F. Charo. (2007). Critical roles for CCR2 and MCP-3 in monocyte mobilization from bone marrow and recruitment to inflammatory sites. *J Clin Invest*, 117(4), 902-909. doi:10.1172/JCI29919

Unger, M. M., J. Spiegel, K. U. Dillmann, D. Grundmann, H. Philippeit, J. Burmann, K. Fassbender, A. Schwiertz and K. H. Schafer. (2016). Short chain fatty acids and gut microbiota differ between patients with Parkinson's disease and age-matched controls. *Parkinsonism Relat Disord*, 32, 66-72. doi:10.1016/j.parkreldis.2016.08.019

Uschold-Schmidt, N., K. D. Nyuyki, A. M. Fuchsl, I. D. Neumann and S. O. Reber. (2012). Chronic psychosocial stress results in sensitization of the HPA axis to acute heterotypic stressors despite a reduction of adrenal in vitro ACTH responsiveness. *Psychoneuroendocrinology*, 37(10), 1676-1687. doi:10.1016/j.psyneuen.2012.02.015

Uschold-Schmidt, N., D. Peterlik, A. M. Fuchsl and S. O. Reber. (2013). HPA axis changes during the initial phase of psychosocial stressor exposure in male mice. *J Endocrinol*, 218(2), 193-203. doi:10.1530/JOE-13-0027

van de Wouw, M., M. Boehme, J. M. Lyte, N. Wiley, C. Strain, O. O'Sullivan, G. Clarke, C. Stanton, T. G. Dinan and J. F. Cryan. (2018). Short-chain fatty acids: microbial metabolites that alleviate stress-induced brain-gut axis alterations. *J Physiol*. doi:10.1113/JP276431

van de Wouw, M., H. Schellekens, T. G. Dinan and J. F. Cryan. (2017). Microbiota-Gut-Brain Axis: Modulator of Host Metabolism and Appetite. *J Nutr*, 147(5), 727-745. doi:10.3945/jn.116.240481

van der Lugt, B., F. Rusli, C. Lute, A. Lamprakis, E. Salazar, M. V. Boekschoten, G. J. Hooiveld, M. Muller, J. Vervoort, S. Kersten, C. Belzer, D. E. G. Kok and W. T. Steegenga. (2018). Integrative analysis of gut microbiota composition, host colonic gene expression and intraluminal metabolites in aging C57BL/6J mice. *Aging (Albany NY)*, 10(5), 930-950. doi:10.18632/aging.101439

van Duin, D., S. Mohanty, V. Thomas, S. Ginter, R. R. Montgomery, E. Fikrig, H. G. Allore, R. Medzhitov and A. C. Shaw. (2007). Age-associated defect in human TLR-1/2 function. *J Immunol*, 178(2), 970-975.

van Kessel, S. P., A. K. Frye, A. O. El-Gendy, M. Castejon, A. Keshavarzian, G. van Dijk and S. El Aidy. (2019). Gut bacterial tyrosine decarboxylases restrict levels of levodopa in the treatment of Parkinson's disease. *Nat Commun*, 10(1), 310. doi:10.1038/s41467-019-08294-y

Varga, G., J. Ehrchen, A. Brockhausen, T. Weinhage, N. Nippe, M. Belz, A. Tsianakas, M. Ross, D. Bettenworth, T. Spieker, M. Wolf, R. Lippe, K. Tenbrock, P. J. Leenen, J. Roth and C. Sunderkotter. (2014). Immune suppression via glucocorticoid-stimulated monocytes: a novel mechanism to cope with inflammation. *J Immunol*, 193(3), 1090-1099. doi:10.4049/jimmunol.1300891

Varga, G., J. Ehrchen, A. Tsianakas, K. Tenbrock, A. Rattenholl, S. Seeliger, M. Mack, J. Roth and C. Sunderkoetter. (2008). Glucocorticoids induce an activated, anti-inflammatory monocyte subset in mice that resembles myeloid-derived suppressor cells. *J Leukoc Biol*, 84(3), 644-650. doi:10.1189/jlb.1107768

Varol, C., L. Landsman, D. K. Fogg, L. Greenshtein, B. Gildor, R. Margalit, V. Kalchenko, F. Geissmann and S. Jung. (2007). Monocytes give rise to mucosal, but not splenic, conventional dendritic cells. *J Exp Med*, 204(1), 171-180. doi:10.1084/jem.20061011

Varol, C., A. Vallon-Eberhard, E. Elinav, T. Ayche, Y. Shapira, H. Luche, H. J. Fehling, W. D. Hardt, G. Shakhbar and S. Jung. (2009). Intestinal lamina propria dendritic cell subsets have different origin and functions. *Immunity*, 31(3), 502-512. doi:10.1016/j.immuni.2009.06.025

Veenstra, M., D. W. Williams, T. M. Calderon, K. Anastos, S. Morgello and J. W. Berman. (2017). Frontline Science: CXCR7 mediates CD14(+)CD16(+) monocyte transmigration across the blood brain barrier: a potential therapeutic target for NeuroAIDS. *J Leukoc Biol*, 102(5), 1173-1185. doi:10.1189/jlb.3HI0517-167R

Vitlic, A., J. M. Lord and A. C. Phillips. (2014). Stress, ageing and their influence on functional, cellular and molecular aspects of the immune system. *Age (Dordr)*, 36(3), 9631. doi:10.1007/s11357-014-9631-6

Vodicka, M., P. Ergang, T. Hrnecir, A. Mikulecka, P. Kvapilova, K. Vagnerova, B. Sestakova, A. Fajstova, P. Hermanova, T. Hudcovic, H. Kozakova and J. Pacha. (2018). Microbiota affects the expression of genes involved in HPA axis regulation and local metabolism of glucocorticoids in chronic psychosocial stress. *Brain Behav Immun*. doi:10.1016/j.bbi.2018.07.007

Vogt, N. M., R. L. Kerby, K. A. Dill-McFarland, S. J. Harding, A. P. Merluzzi, S. C. Johnson, C. M. Carlsson, S. Asthana, H. Zetterberg, K. Blennow, B. B. Bendlin and F. E. Rey. (2017). Gut microbiome alterations in Alzheimer's disease. *Sci Rep*, 7(1), 13537. doi:10.1038/s41598-017-13601-y

Wagner, K. V., X. D. Wang, C. Liebl, S. H. Scharf, M. B. Muller and M. V. Schmidt. (2011). Pituitary glucocorticoid receptor deletion reduces vulnerability to chronic stress. *Psychoneuroendocrinology*, 36(4), 579-587. doi:10.1016/j.psyneuen.2010.09.007

Wattanant, S., D. Tornero, N. Graubardt, T. Memanishvili, E. Monni, J. Tatarishvili, G. Miskinyte, R. Ge, H. Ahlenius, O. Lindvall, M. Schwartz and Z. Kokaia. (2016). Monocyte-Derived Macrophages

Contribute to Spontaneous Long-Term Functional Recovery after Stroke in Mice. *J Neurosci*, 36(15), 4182-4195. doi:10.1523/JNEUROSCI.4317-15.2016

Weise, G., C. Posel, K. Moller, A. Kranz, N. Didwischus, J. Boltze and D. C. Wagner. (2017). High-dosage granulocyte colony stimulating factor treatment alters monocyte trafficking to the brain after experimental stroke. *Brain Behav Immun*, 60, 15-26. doi:10.1016/j.bbi.2016.08.008

White, K. A., S. R. Hutton, J. M. Weimer and P. A. Sheridan. (2016). Diet-induced obesity prolongs neuroinflammation and recruits CCR2(+) monocytes to the brain following herpes simplex virus (HSV)-1 latency in mice. *Brain Behav Immun*, 57, 68-78. doi:10.1016/j.bbi.2016.06.007

Wijeyekoon, R. S., D. Kronenberg-Versteeg, K. M. Scott, S. Hayat, J. L. Jones, M. R. Clatworthy, R. A. Floto, R. A. Barker and C. H. Williams-Gray. (2018). Monocyte Function in Parkinson's Disease and the Impact of Autologous Serum on Phagocytosis. *Front Neurol*, 9, 870. doi:10.3389/fneur.2018.00870

Winek, K., O. Engel, P. Koduah, M. M. Heimesaat, A. Fischer, S. Bereswill, C. Dames, O. Kershaw, A. D. Gruber, C. Curato, N. Oyama, C. Meisel, A. Meisel and U. Dirnagl. (2016). Depletion of Cultivable Gut Microbiota by Broad-Spectrum Antibiotic Pretreatment Worsens Outcome After Murine Stroke. *Stroke*, 47(5), 1354-1363. doi:10.1161/STROKEAHA.115.011800

Wohleb, E. S. (2016). Neuron-Microglia Interactions in Mental Health Disorders: "For Better, and For Worse". *Front Immunol*, 7, 544. doi:10.3389/fimmu.2016.00544

Wohleb, E. S., D. B. McKim, D. T. Shea, N. D. Powell, A. J. Tarr, J. F. Sheridan and J. P. Godbout. (2014). Re-establishment of anxiety in stress-sensitized mice is caused by monocyte trafficking from the spleen to the brain. *Biol Psychiatry*, 75(12), 970-981. doi:10.1016/j.biopsych.2013.11.029

Wohleb, E. S., N. D. Powell, J. P. Godbout and J. F. Sheridan. (2013). Stress-induced recruitment of bone marrow-derived monocytes to the brain promotes anxiety-like behavior. *J Neurosci*, 33(34), 13820-13833. doi:10.1523/JNEUROSCI.1671-13.2013

Woo, V. and T. Alenghat. (2017). Host-microbiota interactions: epigenomic regulation. *Curr Opin Immunol*, 44, 52-60. doi:10.1016/j.coi.2016.12.001

Xia, G. H., C. You, X. X. Gao, X. L. Zeng, J. J. Zhu, K. Y. Xu, C. H. Tan, R. T. Xu, Q. H. Wu, H. W. Zhou, Y. He and J. Yin. (2019). Stroke Dysbiosis Index (SDI) in Gut Microbiome Are Associated With Brain Injury and Prognosis of Stroke. *Front Neurol*, 10, 397. doi:10.3389/fneur.2019.00397

Yano, J. M., K. Yu, G. P. Donaldson, G. G. Shastri, P. Ann, L. Ma, C. R. Nagler, R. F. Ismagilov, S. K. Mazmanian and E. Y. Hsiao. (2015). Indigenous bacteria from the gut microbiota regulate host serotonin biosynthesis. *Cell*, 161(2), 264-276. doi:10.1016/j.cell.2015.02.047

Yeager, M. P., P. A. Pioli, J. Collins, F. Barr, S. Metzler, B. D. Sites and P. M. Guyre. (2016). Glucocorticoids enhance the in vivo migratory response of human monocytes. *Brain Behav Immun*, 54, 86-94. doi:10.1016/j.bbi.2016.01.004

Yin, W., N. R. Gallagher, C. M. Sawicki, D. B. McKim, J. P. Godbout and J. F. Sheridan. (2019). Repeated social defeat in female mice induces anxiety-like behavior associated with enhanced myelopoiesis and increased monocyte accumulation in the brain. *Brain Behav Immun*. doi:10.1016/j.bbi.2019.01.015

Yona, S., K. W. Kim, Y. Wolf, A. Mildner, D. Varol, M. Breker, D. Strauss-Ayali, S. Viukov, M. Guillemins, A. Misharin, D. A. Hume, H. Perlman, B. Malissen, E. Zelzer and S. Jung. (2013). Fate mapping reveals origins and dynamics of monocytes and tissue macrophages under homeostasis. *Immunity*, 38(1), 79-91. doi:10.1016/j.immuni.2012.12.001

Zanier, E. R., F. Marchesi, F. Ortolano, C. Perego, M. Arabian, T. Zoerle, E. Sammali, F. Pischituta and M. G. De Simoni. (2016). Fractalkine Receptor Deficiency Is Associated with Early Protection but Late Worsening of Outcome following Brain Trauma in Mice. *J Neurotrauma*, 33(11), 1060-1072. doi:10.1089/neu.2015.4041

Zeng, Q., G. Junli, X. Liu, C. Chen, X. Sun, H. Li, Y. Zhou, C. Cui, Y. Wang, Y. Yang, A. Wu, Y. Shu, X. Hu, Z. Lu, S. G. Zheng, W. Qiu and Y. Lu. (2019). Gut dysbiosis and lack of short chain fatty acids in a Chinese cohort of patients with multiple sclerosis. *Neurochem Int*, 129, 104468. doi:10.1016/j.neuint.2019.104468

Zeraati, M., M. Enayati, L. Kafami, S. H. Shahidi and A. A. Salari. (2019). Gut microbiota depletion from early adolescence alters adult immunological and neurobehavioral responses in a mouse model of multiple sclerosis. *Neuropharmacology*, 157, 107685. doi:10.1016/j.neuropharm.2019.107685

Zhang, R., R. G. Miller, C. Madison, X. Jin, R. Honrada, W. Harris, J. Katz, D. A. Forsheew and M. S. McGrath. (2013). Systemic immune system alterations in early stages of Alzheimer's disease. *J Neuroimmunol*, 256(1-2), 38-42. doi:10.1016/j.jneuroim.2013.01.002

Zheng, X., S. Ma, A. Kang, M. Wu, L. Wang, Q. Wang, G. Wang and H. Hao. (2016). Chemical dampening of Ly6C(hi) monocytes in the periphery produces anti-depressant effects in mice. *Sci Rep*, 6, 19406. doi:10.1038/srep19406

Zhou, J., L. Tang, C. L. Shen and J. S. Wang. (2018). Green tea polyphenols modify gut-microbiota dependent metabolisms of energy, bile constituents and micronutrients in female Sprague-Dawley rats. *J Nutr Biochem*, 61, 68-81. doi:10.1016/j.jnutbio.2018.07.018

Zhuang, Z. Q., L. L. Shen, W. W. Li, X. Fu, F. Zeng, L. Gui, Y. Lu, M. Cai, C. Zhu, Y. L. Tan, P. Zheng, H. Y. Li, J. Zhu, H. D. Zhou, X. L. Bu and Y. J. Wang. (2018). Gut Microbiota is Altered in Patients with Alzheimer's Disease. *J Alzheimers Dis*, 63(4), 1337-1346. doi:10.3233/JAD-180176

Ziegler-Heitbrock, L., P. Ancuta, S. Crowe, M. Dalod, V. Grau, D. N. Hart, P. J. Leenen, Y. J. Liu, G. MacPherson, G. J. Randolph, J. Scherberich, J. Schmitz, K. Shortman, S. Sozzani, H. Strobl, M. Zembala, J. M. Austyn and M. B. Lutz. (2010). Nomenclature of monocytes and dendritic cells in blood. *Blood*, 116(16), e74-80. doi:10.1182/blood-2010-02-258558

Ziegler-Heitbrock, L. and T. P. Hofer. (2013). Toward a refined definition of monocyte subsets. *Front Immunol*, 4, 23. doi:10.3389/fimmu.2013.00023